

Huang, Y. et al. 2011

A New Variety of Hemiboea (Gesneriaceae) from Limestone Areas in Guangxi, China.

Taiwania 56(3): 240-243.

REFNO: 3759

KEYWORDS:

Guangxi, Hemiboea follicularis var. retroflexa, limestone flora, new variety



A New Variety of *Hemiboea* (Gesneriaceae) from Limestone Areas in Guangxi, China

Yu-Song Huang⁽¹⁾, Wei-Bin Xu^(1,2), Ri-Cheng Peng^(1,3) and Yan Liu^(1*)

1. Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and the Chinese Academy of Sciences, Guilin 541006, China.

2. College of Forestry, South China Agricultural University, Guangdong, Guangzhou 510642, China.

3. College of Life Sciences, Guangxi Normal University, Guilin, 541004, China.

* Corresponding author. Email: gxibly@163.com

(Manuscript received 11 March 2011; accepted 29 April 2011)

ABSTRACT: *Hemiboea follicularis* C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang, a new variety of Gesneriaceae from limestone areas in Guangxi, China, is described and illustrated. The new variety is similar to *Hemiboea follicularis* C. B. Clarke var. *follicularis*, but differs in the stem quadrangular, the corolla yellow green, with purple stria or macula, inside with half-ring of hairs, labels retroflexed.

KEY WORDS: Guangxi; *Hemiboea follicularis* var. *retroflexa*; limestone flora; new variety.

INTRODUCTION

In the course of floristic investigation in limestone areas of Guangxi, China, we discovered a rare plant of *Hemiboea* (Gesneriaceae). After comparing with all described species of *Hemiboea* (Li, 1983, 1987; Wang et al., 1990; Wei et al., 1995; Weitzman et al., 1997; Wang et al., 1998; Li and Liu, 2004; Li and Wang, 2004; Xu et al., 2010), we found it is different from all the known species of *Hemiboea* except *Hemiboea follicularis*, but compared with *Hemiboea follicularis*, it also differs in the stem quadrangular, the corolla yellow green, with purple stria or macula, inside with half-ring of hairs, labels retroflexed. So we confirmed it is a new variety and described as below.

TAXONOMIC TREATMENTS

Hemiboea follicularis C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang, var. nov. Figs. 1 & 2 A-D

H. follicularis C. B. Clarke var. *follicularis similis*, a qua differt caule quadrangulo, corolla luteo-viridi, purpuleo-striata vel maculata, intus pilis semiannualis ornata, labellis reflexis.

Type: China, Guangxi: Guilin, Yanshan, Guilin Botanical Garden, introduced from Huanjiang County by Yan Liu, alt. 170 m, 26 July 2007, W. B. Xu 07066 (holotype IBK).

Herbs perennial. Stems 20-40 cm in height, quadrangular, glabrous, simple or branched, with 4-8 nodes. Leaves opposite, with leaf blade elliptic or ovate lanceolate, 3-15 × 1.5-9 cm; base cuneate, usually

asymmetric; margin nearly entire or weakly sinuous; apex acuminate or acute; adaxial blade green, glabrous; abaxial blade green-white, veins with sparsely pubescence; lateral veins 5-8 on each side of midrib; petiole 1-8 cm long. Cymes 5-10 or more flowers, axillary or pseudoterminal, peduncle 3-5 cm long, glabrous. Involucre 1.5-2.5 cm in diam., globular; apex white, membranous, glabrous, bowl-shaped when opened. Calyx 1-1.5 cm long, 5-lobed from above middle, white, glabrous. Corolla yellow green, with purple stria or macula, 1.2-1.8 cm long, outside glabrous, corolla tube 1-1.5 cm long, inside densely short glandular-hairy and puberulent, with half-turn of hairs adnate to 4-6 mm above the corolla base, adaxial lip 2-3 mm long, 2-lobed, lobe hemicycle, retroflexed, abaxial lip 2.5-3.5 mm long, 3-lobed, lobe oval, retroflexed. Stamen 2; filaments linear, 1-1.5 mm long, adnate to 4-6 mm above the corolla base; anthers elliptic, 2 mm long, connate completely at the venter; staminodes 2, ca. 1.5 mm long; apex capitate. Disc annular, ca. 1.2 mm high. Pistil 6-8 mm long; ovary linear, glabrous; stigma capitellate; apex densely spheroidal murication. Capsule unknown.

Ecology: On shaded rocky limestone slopes in broadleaved forests, ca. 200-300 m elevation.

Phenology: Flowering June to July.

Etymology: The specific epithet is derived from the shape of corolla lobes whose adaxial and abaxial lip retroflexed.

IUCN Red List Category: The species are only found from Longzhou and Huanjiang County, Guangxi, China at now. Current information shows that the population size of this new species is fewer than 250 mature individuals. We therefore assess *Hemiboea follicularis*




Fig. 1. *Hemiboea follicularis* C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang A: Habit. B: Flower. C: Calyx. D: Corolla opened, showing stamens and staminodes. E: Pistil. F: Stigma. Drawn by S. Q. He from W. B. Xu 07066 (IBK).

var. *retroflexa* as endangered degree (EN) using the IUCN categories and criteria (IUCN 2001).

Note: *Hemiboea follicularis* C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang is closely related to *Hemiboea follicularis* C. B. Clarke var. *follicularis* (Fig. 2 E & F), but differs in the stem quadrangular, the corolla yellow green, with purple stria or macula, inside with half-ring of hairs, labels retroflexed. A comparison of the salient characters of *Hemiboea follicularis* var. *retroflexa* and *H. follicularis* var. *follicularis* is shown in

Table 1.

Additional specimens examined: CHINA. Guangxi: Longzhou County, Chunxiu natural reserve, on shaded rocky limestone slopes in broadleaved forests, rare, alt. 250 m, 23 July 2009, R. H. Jiang et al. H09603 (IBK).

ACKNOWLEDGEMENTS

We are grateful to Prof. Fa-Nan Wei (IBK) for the Latin diagnosis, to Mr. Shun-Qing He (IBK) for preparing the handsome illustration. We also thank the staff of Chunxiu




Fig. 2. A-D: *Hemiboea follicularis* C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang. A: Habit. B: Involucre. C: Lateral flower. D: Frontal flower. E, F: *Hemiboea follicularis* C. B. Clarke var. *follicularis*. E: Habit. F: Flower.

Table 1. Comparison of salient characters of *Hemiboea follicularis* var. *retroflexa* and *H. follicularis* var. *follicularis*

	<i>Hemiboea follicularis</i> var. <i>retroflexa</i>	<i>H. follicularis</i> var. <i>follicularis</i>
Stem	quadrangular	rounded
Corolla	yellow green, with purple stria or macula, inside with half-ring of hairs	white, inside with a ring of hairs
Labels	retroflexed, abaxial lip oval	not retroflexed, abaxial lip half-rounded
Cymes	5-10 or more flowers, axillary or pseudo terminal	7-20 or more flowers, pseudoterminal
Anther	elliptic, 2 mm long	rounded, 1-2 mm long



natural reserve of Longzhou for field assistance. This study was supported by Knowledge Innovation Project of the Chinese Academy of Sciences, Grant No. KSCX2-YW-Z-0912 to Yan Liu (IBK), Fundamental Research Fund of Guangxi Institute of Botany (GZY 09004) to Wei-Bin Xu (IBK) and Scientific Funded Projects of Guangxi (GKZ0991226).

LITERATURE CITED

- IUCN. 2001. IUCN Red List Categories and Criteria (Version 3.1). IUCN Publication Service Unit, Gland, Switzerland and Cambridge.
- Li, Z.-Y. 1983. Taxa nova *Hemiboeae* (Gesneriaceae). Acta Phytotax. Sin. **21**: 194-203. [in Chinese]
- Li, Z.-Y. 1987. A study of the genus *Hemiboeae* (Gesneriaceae). Acta Phytotax. Sin. **25**: 220-230. [in Chinese]
- Li, Z.-Y. and Y. Liu. 2004. *Hemiboea rubribracteata* Z. Y. Li & Yan Liu, a new species of *Hemiboea* (Gesneriaceae) from Guangxi, China. Acta Phytotax. Sin. **42**: 537-540. [in Chinese]
- Li, Z.-Y. and Y.-Z. Wang. 2004. Plants of Gesneriaceae in China. Henan Science and Technology Publishing House, pp. 123-153. [in Chinese]
- Wang, W.-T., K.-Y. Pan, Z.-Y. Zhang, Z.-Y. Li, D. D. Tai and W.-C. Yin. 1990. Gesneriaceae. In: Flora Reipubl. Pop. Sin. **69**: 125-581, Science Press, Beijing, China. [in Chinese]
- Wang, W.-T., K.-Y. Pan, Z.-Y. Li, A. L. Weitzman and L. E. Skog. 1998. Gesneriaceae. In: Wu, Z. H. and P. H. Raven (eds.), Flora of China. **18**: 244-401, Science Press, Miss. Bot. Gard. Press.
- Wei, Y.-G. and H.-Q. Wen. 1995. Two new species from Guangxi. Guihaia **15**: 216-219. [in Chinese]
- Weitzman, A. L., L. E. Skog, W.-T. Wang, K.-Y. Pan and Z.-Y. Li. 1997. New taxa, new combination, and notes on Chinese Gesneriaceae. Novon **7**: 423-435.
- Xu, W.-B., W.-H. Wu, D.-X. Nong and Y. Liu. 2010. *Hemiboea purpurea* sp. nov. (Gesneriaceae) from a limestone area in Guangxi, China. Nord. J. Bot. **28**: 313-315.

中國廣西石灰岩地區半蒴苣苔屬（苦苣苔科, Gesneriaceae）一新變種

黃俞淞⁽¹⁾、許為斌^(1,2)、彭日成^(1,3)、劉演^(1*)

1. 廣西壯族自治區 中國科學院廣西植物研究所，桂林 541006，中國。

2. 華南農業大學林學院，廣州 510642，中國。

3. 廣西師範大學生命科學學院，桂林 541004，中國。

* 通信作者。Email: gxibly@163.com

(收稿日期：2011年3月11日；接受日期：2011年4月29日)

摘要：描述了中國廣西石灰岩地區苦苣苔科（Gesneriaceae）一新變種－卷瓣半蒴苣苔（*Hemiboea follicularis* C. B. Clarke var. *retroflexa* Yan Liu & Y. S. Huang）。該變種與原變種－華南半蒴苣苔（*Hemiboea follicularis* C. B. Clarke var. *follicularis*）相近，不同在於莖四棱形，花冠黃綠色，具紫色條紋或斑點，花冠內部只具半圈毛環，唇瓣反卷，可以區別。

關鍵詞：廣西、卷瓣半蒴苣苔、石灰岩植物區系、新變種。