

**Wu, W.H., Xu, Nong, & Liu 2011**

**Chirita ningmingensis (Gesneriaceae), a new species from Guangxi, China.**

**Ann. Bot. Fenn. 48(5): 422-424.**

**REFNO: 3838**

**KEYWORDS:**

**China, Chirita, Guangxi, new species**

## ***Chirita Ningmingensis* (Gesneriaceae), a New Species from Guangxi, China**

Author(s): Wang-Hui Wu, Wei-Bin Xu, Dong-Xing Nong & Yan Liu

Source: *Annales Botanici Fennici*, 48(5):422-424. 2011.

Published By: Finnish Zoological and Botanical Publishing Board

URL: <http://www.bioone.org/doi/full/10.5735/085.048.0505>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

## *Chirita ningmingensis* (Gesneriaceae), a new species from Guangxi, China

Wang-Hui Wu<sup>1,2</sup>, Wei-Bin Xu<sup>1</sup>, Dong-Xing Nong<sup>1,2</sup> & Yan Liu<sup>1,\*</sup>

<sup>1</sup> Guangxi Institute of Botany, Guangxi Zhuangzu Autonomous Region and the Chinese Academy of Sciences, Guilin 541006, Guangxi, China (\*corresponding author's e-mail: gxibly@163.com)

<sup>2</sup> College of Life Sciences, Guangxi Normal University, Guilin 541004, Guangxi, China

Received 15 Apr. 2010, revised version received 19 May 2010, accepted 10 June 2010

Wu, W. H., Xu, W. B., Nong, D. X. & Liu, Y. 2011: *Chirita ningmingensis* (Gesneriaceae), a new species from Guangxi, China. — *Ann. Bot. Fennici* 48: 422–424.

*Chirita ningmingensis* Yan Liu & W.H. Wu, a new species of Gesneriaceae from Guangxi, China, is described and illustrated. It is similar to *C. baishouensis* but differs in the peduncle and pedicel glandular-pubescent, bracts broadly lanceolate, the corolla 3–3.5 cm long, outside glandular-pubescent, filaments ca. 6 mm long, anthers glabrous, staminodes 3, the pistil glandular-pubescent, stigma cordiform, and apex retuse.

There are more than 60 species of *Chirita* (Gesneriaceae) in Guangxi (Qin & Liu 2010), including some recently described ones (Liu & Wei 2004, Li *et al.* 2006, Wei *et al.* 2007, Li & Wang 2008, Pan *et al.* 2009). In the recent years, we carried out some floristic surveys in the limestone caves resulting in the discovery of three new species from Guangxi (Liu *et al.* 2007, Xu *et al.* 2008, 2009). In this paper, another new species of *Chirita* from a limestone cave in the karst areas of southwestern Guangxi is described.


***Chirita ningmingensis*** Yan Liu & W.H. Wu, *sp. nova* (Fig. 1)

*Species nova C. baishouensi similis, sed pedunculis pedicellisque glanduloso-pubescentibus, bracteis late lanceolatis, corolla 3–3.5 cm longa, extus glanduloso-pubescentibus, filamentis circ. 6 mm longis, antheris glabris, staminodiis 3, pistillo glanduloso-pubescenti, stigmatibus cordato, apice retuso differt.*

TYPE: China. Guangxi, Ningming County, Longgang National Nature Reserve, at the entrance of a limestone cave, rare, alt. 250 m, 18.VIII.2002 Yan Liu L0718 (holotype IBK; isotype PE). — PARATYPE: China. Guangxi, Ningming County, Longgang National Nature Reserve, rare, alt. 250 m, 23.VIII.2009 Wang-Hui Wu W09155 (IBK).

ETYMOLOGY. The specific epithet is derived from the type locality, Longgang National Nature Reserve, Ningming County, Guangxi.

Herbs perennial. Rhizome subterete, 2–4 cm long, 0.8–1.2 cm wide. Leaves petiolate; petiole applanate, 3–8 cm long, 3–4 mm wide, appressed pubescent; leaf blade herbaceous, ovate or broadly elliptic to subrounded, 4–9 × 3–7 cm, appressed pubescent on both surfaces, apex acute to obtuse, base cuneate, sometimes inequilateral, margin entire, lateral veins 3–5 on each side. Cymes 5–10, axillary, 1–2-branched, 8–20-flowered; peduncle 4–11 cm long, 1.5–3 mm wide, glandular-pubescent; bracts broadly lanceolate, 0.8–2.5 × 0.4–1.1 cm, apex acute, margin entire, appressed pubescent on both surfaces, pedicel 1–3 cm long, ca. 1.2 mm wide, glandular-pubes-



**Fig. 1.** *Chirita ningmingensis* (from the holotype, drawn by S. Q. He). — **A:** Habit. — **B:** Corolla opened with stamens and staminodes. — **C:** Calyx and pistil. — **D:** Stigma.

cent. Calyx 5-parted nearly to base, 7–9 × ca. 1 mm, lobes linear-lanceolate, outside glandular-pubescent, inside glabrous, apex acuminate, margin entire. Corolla pale purple-red, 3–3.5 cm long, outside glandular-pubescent, inside sparsely glandular-pubescent, tube 1.8–2 cm long, 8–10 mm in diam. at mouth, ca. 4 mm in diam. at base; limb distinctly 2-lipped, adaxial lip 2-lobed to over middle, lobes oblong, 3–4 × ca. 6 mm, abaxial lip 3-lobed to near middle, lobes oblong, 5–6 × ca. 5 mm, with 2 yellow stripes inside. Stamens 2, adnate ca. 1.5 cm above the corolla base; filaments ca. 6 mm long, glandular-pubescent, strongly geniculate near middle, anthers ca. 2 mm long, glabrous.

Staminodes 3, lateral ones ca. 3.5 mm long, apex capitate, sparsely pubescent, adnate to 1.1 cm above corolla tube base, middle one 0.5–1 mm long, adnate to 5 mm above corolla tube base. Disc ca. 1.5 mm high, margin repand, glabrous. Pistil 2.2–2.5 cm long, ovary 1.2–1.5 cm, long, 1.5 mm wide, glandular-pubescent; style 8–10 mm long, 1.5 mm wide, glandular-pubescent; stigma cordiform, apex retuse, 1.3 mm wide. Capsule not observed. Flowering from August to September.

*Chirita ningmingensis* is known only from one population at the entrance of a limestone cave in the Longgang National Nature Reserve, Ningming County, Guangxi. It is similar to *C.*

**Table 1.** Morphological comparison of *Chirita ningmingensis* and *C. baishouensis*.

Characters	<i>Chirita ningmingensis</i>	<i>C. baishouensis</i>
Peduncle and pedicel	glandular-pubescent	pubescent
Bracts	broadly lanceolate	narrowly lanceolate
Calyx	lobes outside glandular-pubescent, inside glabrous, margin entire	lobes appressed pubescent on both surfaces, margin sparsely denticulate
Corolla	3–3.5 cm long, outside glandular-pubescent, with 2 yellow stripes inside	(3.7–)4.5–6.8 cm long, outside pubescent; with 4 purple stripes inside
Filaments	ca. 6 mm long, glandular-pubescent	1–1.4 cm long, glabrous or lower part sparsely glandular-pubescent
Anthers	ca. 2 mm long, glabrous	(1.5–)5 mm long, densely barbate on backside
Staminodes	3, sparsely pubescent	2, sparsely glandular-pubescent
Pistil	glandular-pubescent	densely appressed pubescent
stigma	cordiform, apex retuse	2-lobed, lobes narrowly triangular
Flowering	August to September	March to April

*baishouensis* (Wei et al. 2000), but it can be distinguished e.g. by the peduncle and pedicel glandular-pubescent, bracts broadly lanceolate, the corolla 3–3.5 cm long, outside glandular-pubescent, filaments ca. 6 mm long, anthers glabrous, staminodes 3, the pistil glandular-pubescent, stigma cordiform, and apex retuse (see Table 1).

## Acknowledgements

We are grateful to Prof. Fa-nan Wei (IBK) for the Latin diagnosis. We also thank Mr. Shun-Qing He (IBK) for preparing the illustration. This study was supported by Knowledge Innovation Project of the Chinese Academy of Sciences, Grant No. KSCX2-YW-Z-0912 to Yan Liu (IBK), and Fundamental Research Fund of Guangxi Institute of Botany (GZY 09004) to Wei-bin Xu (IBK).

## References

- Li, Z. Y., Li, Y. B. & Xing, Q. 2006: *Chirita tribracteata* var. *zhuana* Z. Y. Li, Q. Xing & Y. B. Li (Gesneriaceae), a new variety from Guangxi, China. — *Acta Phytotaxonomica Sinica* 44: 649–650.
- Li, J. M. & Wang, Y. Z. 2008: *Chirita longicalyx* (Gesneriaceae), a new species from Guangxi, China. — *Annales Botanici Fennici* 45: 212–214.
- Liu, Y. & Wei, Y. G. 2004: [*Chirita lutea* Yan Liu et Y. G. Wei, a new species of Gesneriaceae from Guangxi, China.] — *Journal of Wuhan Botanical Research* 22: 391–393. [In Chinese].
- Liu, Y., Wei, Y. G. & Tang, S. C. 2006: *Chiritopsis lingchuanensis* Yan Liu et Y. G. Wei, a new species of Gesneriaceae from Guangxi, China. — *Acta Phytotaxonomica Sinica* 44: 340–344.
- Pan, B., Mo, D., Xu, W. B. & Zhou, T. J. 2009: *Chirita pseudoheterotricha* T. J. Zhou, B. Pan & W. B. Xu, A New Species of Gesneriaceae from Limestone Areas in Guangxi, China. — *Journal of Tropical and Subtropical Botany* 17: 596–599.
- Qin, H. N. & Liu, Y. 2010: [*A checklist of vascular plants of Guangxi*]. — Sci. Press, Beijing. [In Chinese].
- Wei, Y. G., Pan, B. & Tang, W. X. 2007: *Chirita guihaiensis* sp. nov. (Gesneriaceae) from Guangxi, China. — *Nordic Journal of Botany* 25: 296–298.
- Wei, Y. G., Wen, H. Q. & Zhong, S. H. 2000: New materials of Gesneriaceae from Guangxi, China. — *Acta Phytotaxonomica Sinica* 38: 297–301.
- Xu, W. B., Gao, H. S. & Liu, Y. 2008: *Lagarosolen jingxiensis* Yan Liu, H. S. Gao & W. B. Xu, a New Species of Gesneriaceae from Guangxi, China. — *Journal of Tropical and Subtropical Botany* 16: 274–276.
- Xu, W. B., Liu, Y. & Gao, H. S. 2009: *Chiritopsis jingxiensis*, a new species of Gesneriaceae in karst cave from Guangxi, China. — *Novon* 19: 559–561.