

**A new species of Solenophora (Gesneriaceae) from
Southeast Veracruz, Mexico.**

Novon 7: 281-284.

REFNO: 2699


KEYWORDS:

Mexico, Solenophora



PK
1
N946
BOT

Missouri
Botanical
Garden



Novon

A Journal for Botanical Nomenclature

VOLUME 7 NUMBER 3

1997

A New Species of *Solenophora* (Gesneriaceae) from Southeast Veracruz, Mexico

Angélica Ramírez-Roa* and Guillermo Ibarra-Manríquez**

*Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 70-233, Deleg. Coyoacán, 04510 México, D.F., Mexico.

**Dirección actual: Departamento de Ecología de los Recursos Naturales, Instituto de Ecología, Universidad Nacional Autónoma de México, Apartado Postal 27-3, Xangari 58089, Morelia, Michoacán, Mexico

1979. A new androecial
Bot. Neerl. 28: 349-355.
1 Africa and Madagascar.

ms in African Lorantha-
lm-Nielsen, I. C. Nielsen
ical Forests. Academic

nthaceae, Viscaceae. In:
numération des Plantes à
162-185. Conservatoire

RESUMEN. Se describe e ilustra *Solenophora tuxtliensis* (Gesneriaceae, Gesnerioideae), una nueva especie de Veracruz, México. Las características más evidentes para distinguir esta especie son el cáliz muy largo y tubular, la corola con máculas púrpuras en los lóbulos y una glándula nectarífera bilobulada. *Solenophora tuxtliensis* es conocida solamente de la región de Los Tuxtlas, Veracruz.

ABSTRACT. *Solenophora tuxtliensis* (Gesneriaceae, Gesnerioideae), a new species from Veracruz, Mexico, is described and illustrated. The most prominent features distinguishing this species are the long tubular calyx, and the corolla with dark purple spots on the lobes and only one bilobed nectariferous disc gland. It is known only from the Los Tuxtlas region, Veracruz.

Solenophora Benthham is a neotropical genus in the tribe Gloxinieae (Fritsch) Wiehler (Wiehler, 1983), and has about 15 species occurring from Mexico to Panama (Gibson, 1974; Skog, 1979). Similar to other genera in the family, this genus is poorly collected in Mexico and needs to be revised. About six species in the genus have been collected in Mexico in the states of Chiapas, Oaxaca, Puebla, and Veracruz.

During preparation of a floristic study of the "Los Tuxtlas" Biological Station in Veracruz, Ibarra-Manríquez and Sinaca (1987) collected material of *Solenophora* that corresponded to specimens provisionally named as *Solenophora* "tuxtla" by Dale Denham. An examination of herbarium material from MEXU and US and the literature about this genus revealed to us that the Denham species is worthy of publication.

Solenophora tuxtliensis Ramírez-Roa & Ibarra-Manríquez, sp. nov. TYPE: México. Veracruz: Mpio. San Andrés Tuxtla, Estación de Biología Tropical Los Tuxtlas, Lote 71, 18°34'–36'N. 95°05'–09'W, 400 m, 18 ene. 1991, G. Ibarra & S. Sinaca 3533 (holotype, MEXU; isotypes, BM, ENCB, F, K, LE, MO, US, XAL). Figure 1.

Planta terrestris suffrutescens vel frutex erectus, ramosus; caulis pilosus, folia opposita longipetiolata, herbacea, laxa, pilosa, lamina elliptica; petioli axillari 1–2; calyx 2.6–4.1 cm longus, subcampanulatus flavescens, limbus breviter 5-lobatus, lobi deltoidei serrati; corolla cylindrica, flava, tubus intus purpureo-maculatus, lobi purpureo-maculati; stamina epicorollina, antherae apice connatae; stigma stomatomorphum; ovarium inferum pilosum; discus e glandula bilobata formatus; capsula calyce accrescenti; semina elliptica.

Suffrutescent herbs or shrubs 1.5–2.5 m tall; internodes pubescent, trichomes multicellular, 0.5–1 mm long, more conspicuous in younger parts. Leaves opposite, simple, elliptic, rarely narrowly elliptic or ovate, 3.5–25 × 1.8–14 cm, 1.5–3 times longer than wide, the base oblique, the apex acute or cuspidate, biserrate, membranous, upper surface dark green, pilose, trichomes 1–2 mm long, lower surface light green or frequently reddish, pilose along the costa and secondary veins, 6–9 secondary veins, pinnate; petiole 1.5–13 cm long, sparsely pilose, reddish above. Peduncle reddish, 2.5–7 cm long, pilose; bracts 2, linear, 0.4–1.2 cm long, green, pilose. Flowers showy, 1–2 in each axil; floral tube turbinate, 6–9 × 3–6 mm, red at the base, yellowish at the apex; calyx erect, slightly campanulate, 2.6–4.1 cm long, yellow; tube 1.5–3.3 × 0.8–1.5 cm, reddish tomentose; 5 lobes triangular, 2–8 × 2–8 mm long and wide, margin serrate and pubescent, trichomes ca. 1 mm long; corolla tubular, slightly broader toward the apex, 4.3–6 cm long, yellow, with dark purple spots on the lobes, the spots becoming lines toward inside of the tube; tube of the corolla 3.5–5 cm long, 0.7–1 cm broad at the base, bisacciform dorsally, 1.3–1.6 cm at the throat, the limb 5-lobed, 0.8–1 cm long × 7–8 mm wide, margin fimbriate; 4 stamens diadelphous, included, adnate to the base of the corolla tube, 4.4–4.6 cm long, glabrous, anthers broadly ovate, coherent in a square, each 2–4 mm wide, longitudinally dehiscent, the connective glabrous, staminode ca. 5 mm long, pollen tricolpate, colpi 24–29 μm




Figure 1. *Solenophora tuxtensis* Ramírez-Roa & Ibarra-Manríquez. —a. Habit showing flowers. —b. Leaf blade base, abaxial surface. —c. Flower opened, showing diadelphous stamens, staminode, disc gland, style, and stigma. —d. Close-up of petal lobe with fimbriate margin. —e. Close-up of bilobed disc gland (staminode removed). —f. Flower, lateral view (calyx and corolla removed). Based on Ibarra & Sinaca 3533, MEXU.




Figure 2. A-H. Pollen of *Solenophora tuxtlenis* Ramírez-Roa & Ibarra-Manríquez. A-D. Light microscope photomicrographs of pollen (based on Ibarra & Sinaca 3533, MEXU). (Scale 1 cm = 10 μ m). —A. Meridional view. —B. Meridional view. —C. Polar view. —D. Polar view. E-H. SEM photomicrographs of pollen (based on Ibarra & Sinaca 3533, MEXU). —E. Meridional view showing colpus. —F. Meridional view showing two colpi. —G. Polar view. —H. Close-up of mesocolpium and three colpi. I, J. Photomicrographs of *S. tuxtlenis* seed (based on Ibarra & Sinaca 3533, MEXU). —I. Longitudinal view of seed showing shape, apex, and ornamentation. —J. Epidermal cells of testa on face and apex (description in the text follows Beaufort-Murphy, 1983).

rs. —b. Leaf blade base, style, and stigma. —d. e removed). —f. Flower,

Alonso 94

long, sometimes syncolpate, margins distinct, displaying persistent micro-verrucate colpus membrane (in SEM), exine semi-tectate, 1–1.5 μm thick, sexine micro-reticulate, heterobrochate, lumina $< 1 \mu\text{m}$, diminishing gradually toward poles and apertures (SEM), sexine/nexine ratio 2:1, prolate, polar axis 26–34 μm , equatorial axis 17–28 μm (Fig. 2A–H); nectariferous disc of 1 dorsal gland bilobed; ovary inferior, apex slightly convex, pilose to tomentose at the apex, reddish, rarely oblong, style terete, 4.5–5.5 cm long, pilose, stigma stomatomorphic, with glandular trichomes. Dry capsule surrounded by the accrescent calyx, 0.8–1.8 \times 0.4–0.7 cm, turbinate, reddish tomentose; seeds numerous, ca. 5 mm long, elliptic, dark brown, apex constricted, cell shape narrowly linear-polygonate, with edges elevated, crests discrete, face depressed, edges and faces verrucate (Fig. 2I, J). Chromosome number $n = 10$ (Skog, 1984).

?!
probably
.5 mm

Distribution and habitat. The new species is an uncommon plant known only from three municipal areas of Veracruz, Mexico: San Andrés Tuxtla, Catemaco, and Soteapan. It grows on rocks in small rivers, in primary tropical rainforest or cloud forest, at 200–950 m s.n.m.

Reproduction. Flowering is mainly from January to May, fruiting from March to June. Hummingbirds were seen visiting the flowers.

The epithet "tuxtla" suggested by Denham was written on herbarium specimens and labels of living collections. We have changed it to *tuxtensis* following the Code (Recommendation 60 D.I., Greuter et al., 1994).

Paratypes. MEXICO. Veracruz: Municipio Catemaco, entre ejidos Adolfo Ruiz Cortés y La Perla del San Martín, ca. 15 km al N de Catemaco, 18°31'N, 95°05'W, 950 m, 21 feb. 1972 (fr), *Beaman 5760* (MEXU, US); al N de un poblado a 10 km al SE de Tebanca camino al Bastonal, 19 dic. 1984, *Cedillo 2950* (MEXU); Municipio San Andrés Tuxtla, Estación de Biología Tropical Los Tuxtlas, Lote 71, 18°34'–36'N, 95°05'–09'W, 550 m, 17 feb. 1986 (fl, fr), *Ibarra & Sinaca 2824* (MEXU), 600 m, 1 ago. 1986 (fl, fr), *Ibarra et al. 2975* (MEXU, US); Volcán San Martín Tuxtla, "El Paraje" at edge of small stream, 1500 m, 18 Feb. 1962 (fl), *MacDougall 492* (US); Municipio San Andrés Tuxtla, Estación de Biología Tropical Los Tuxtlas, Lote 71, 18°34'–36'N, 95°05'–09'W, 600 m, 2 abr. 1986 (fl, fr), *Sinaca et al. 521* (MEXU, US); Laguna Azul, 200 m, 20 mar. 1986 (fl, fr), *Sinaca 486* (MEXU, US); Laguna Escondida, 18°35'N, 95°01'W, 200 m, 21 abr. 1985 (fl, fr), *Sinaca & Aparicio 73* (MEXU); Laguna Escondida, 20 abr. 1972 (fr), *Villegas 7* (MEXU, US); Municipio Soteapan, along dirt road 13 km E of Tebanca (13 km E of E side of Lago de Catemaco), 800–950 m, 5 July 1980 (fr), *Nee & Hansen 18768* (MEXU). CULTIVATED: U.S.A. New York: Ithaca, grown in the Cornell University Conservatory, as G-911, 23 June 1966 (fl), *Stone 136* (US).

US

At present it is difficult to relate *Solenophora*

tuxtensis to other species of the genus. It is distinctive by its long tubular calyx, its tubular corolla with dark purple spots on the lobes, and by having only one bilobed disc gland. A key to distinguish *S. tuxtensis* from the other species found in Oaxaca and Veracruz is presented:

- 1a. Calyx nearly truncate, lobes 1–2 mm long.
 - 2a. Calyx tubular, base reddish; corolla lobes undulate *S. maculata* D. Gibson
 - 2b. Calyx campanulate, base greenish; corolla lobes fimbriate *S. insignis* (M. Martens & Galeotti) Hanstein
- 1b. Calyx lobulate, lobes 2–8 mm long.
 - 3a. Leaf margins serrate *S. obscura* Hanstein
 - 3b. Leaf margins biserrate *S. tuxtensis* Ramfrez-Roa & Ibarra-Manríquez

Acknowledgments. The senior author thanks Laurence E. Skog (US) for a careful review of the manuscript, for his help in reviewing the specimens at US, and for sharing with us his broad knowledge of gesneriads. The junior author thanks Santiago Sinaca Colín for assistance in the field. Thanks are also given to Patricia Dávila and the staff at MEXU for facilitating our access to the collections and for computer support. We also thank Fernando Chiang (MEXU), who kindly checked the Latin description and pointed out the change of the name; Socorro Lozano and the staff of the Instituto de Geología, Laboratorio de Palinología (UNAM), for helping with the LM and SEM work and the description of the pollen; Albino Luna (Instituto de Biología, UNAM) for the fine botanical illustration; and Felipe Villegas (Instituto de Biología, UNAM) for his generous assistance in making Figure 2.

Literature Cited

- Beaufort-Murphy, H. T. 1983. The seed surface morphology of the Gesneriaceae utilizing the Scanning Electron Microscope and a new system for diagnosing seed morphology. *Selbyana* 6: 220–422.
- Gibson, D. N. 1974. Gesneriaceae. In: P. C. Standley, L. O. Williams & D. N. Gibson (editors), *Flora of Guatemala*. Fieldiana, Bot. 24(10): 240–313.
- Greuter, W., F. R. Barrie, H. M. Burdet, W. G. Chaloner, V. Demoulin, D. L. Hawksworth, P. M. Jorgensen, D. H. Nicolson, P. C. Silva, P. Treharne & J. McNeill. 1994. *International Code of Botanical Nomenclature* (Tokyo Code). *Regnum Veg.* 131.
- Ibarra-Manríquez, G. & S. Sinaca C. 1987. Lista Florística de la Estación de Biología Tropical Los Tuxtlas, Veracruz. *Listados Florísticos de México VII*. Instituto de Biología, Universidad Nacional Autónoma de México, México, D.F.
- Skog, L. 1979. Gesneriaceae. In: R. E. Woodson & R. W. Schery (editors), *Flora of Panama*. *Ann. Missouri Bot. Gard.* 65: 783–996.
- . 1984. A review of chromosome numbers in the Gesneriaceae. *Selbyana* 7: 252–273.
- Wiehler, H. 1983. A synopsis of the neotropical Gesneriaceae. *Selbyana* 6: 1–219.

ABSTR
annul.
iana,
M. me
from t
briata
dônia,
new a
branch
fimbri

In p
Spreng
herbar
cies c
Rondô
to be c
of gra
rich tr
Five
M. cal
are kn
In d
on ste
collect
and en
additio
the pr
The
and al
Rhipid.
insertio
many,
equal,
gle bu
located
culm.

BRANC
Some
and M.
undesc
the bra
of the