

**Determination of the unequal fate of cotyledons of a one-leaf plant, Monophyllaea.**

**Development 124: 1275-1280.**

**REFNO: 3152**

**KEYWORDS:**

**Cotyledon, Development, Malaysia, Monophyllaea, Morphology, Seedling**

## Determination of the unequal fate of cotyledons of a one-leaf plant, *Monophyllaea*

Hirokazu Tsukaya

Institute of Molecular and Cellular Biosciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113, Japan

(e-mail: h33171@m-unix.cc.u-tokyo.ac.jp)

### SUMMARY

One-leaf plants, belonging to the family Gesneriaceae, were described for the first time more than 150 years ago. One such unusual plant, *Monophyllaea*, has only one leaf at maturity. Only one of the two cotyledons grows continuously, without the formation of true leaves, and this feature, known as anisocotyledonous development, has been repeatedly mentioned in textbooks of plant morphology. However, the mechanism for the determination of the one-leaf phenotype remains to be ascertained.

In this study, meristematic regions were identified, by monitoring DNA synthesis, at the base of both cotyledons just after germination, while no such regions were found in

the shoot apex. Surgical experiments with seedlings and analysis of the anisocotyledonous development revealed that the fate of the cotyledons is determined during their growth. Anisocotyledonous development seems to be the result of competition between the two cotyledons. The mechanism that governs the development of the shoot in the genus *Monophyllaea* is discussed in relation to apical dominance, which is the common mechanism that regulates shoot development in many plants.

Key words: anisocotyledonous development, competitive organogenesis, leaf, meristem, *Monophyllaea*, suppression

### INTRODUCTION

*Monophyllaea* was first described in 1840 as an unusual genus of one-leaf plants (Brown, 1840). As the Latin name indicates, each plant has only one leaf at maturity (Fig. 1F). The single, large leaf (more than 30 cm in length in some species) is derived from one of the two cotyledons (Chifflet, 1909; Oehlkers, 1923; Hill, 1938; Jong and Burt, 1975). One of the cotyledons continues to grow while the other does not. No new organs are formed in the shoot after germination until the appearance of the inflorescences (Weber, 1975, 1976a,b). Thus, this plant lacks true leaves. The unequal development of two cotyledons is known as anisocotyledonous development. This feature is common to all known one-leaf species in the family Gesneriaceae (Hill, 1938), which includes more than 30 species in the genus *Monophyllaea* (Burt, 1978) and several species in the genus *Streptocarpus*. Thus, special terms have been created for the description of the unique organs and tissue. 'Phyllomorph' (Jong and Burt, 1975) is the name given to a unit of mixed stem-leaf nature in one-leaf plants (including species in the genera *Streptocarpus* and *Monophyllaea*). A cotyledonary phyllomorph is composed of a large cotyledon and a 'petiolode', which is the petiole-like organ of the large, continuously growing cotyledon. It seems likely that the phyllomorph is maintained by the activity of three kinds of meristem (Jong and Burt, 1975), namely, the 'basal meristem', which is involved in the continued growth of the lamina; the 'petiolode meristem', which produces a long 'petiolode'; and a 'groove meristem', which is essential for the differentiation

of each inflorescence meristem. As the requirement for these individual terms suggests, the unusual development of *Monophyllaea* involves many issues of interest to plant morphologists (e.g., Wardlaw, 1952; Bell, 1991). However, no details of the determination of the unique one-leaf phenotype of *Monophyllaea* are understood. In this study, re-examination of surgical experiments with seedlings, which were attempted previously by Oehlkers (1923), revealed for the first time that the fate of each cotyledon is determined, after germination, by competition.

### MATERIALS AND METHODS

Seeds of *Monophyllaea horsfieldii* R. Br., collected in Malaysia, were sown on rockwool and plants were grown at 23°C under continuous white light as described by Tsukaya et al. (1994). Seeds were stored at room temperature under dry conditions, without any significant loss of the capacity for germination (approx. >95%). For cultivation on plates, seeds were sterilized in a solution of NaClO (Tsukaya et al., 1991) and sown on the medium described by Okada and Shimura (1992). Microscopic observations were made as described previously, with an optic microscope (Leica DAS Mikroskop DMR; Leica, Wetzlar, FRG) and a scanning electron microscope (JMS-820S; JEOL Ltd., Tokyo, Japan; Tsukaya et al., 1993; Tsuge et al., 1996). For identification of cell nuclei in whole-mounted seedlings, fixed samples were stained with 1 µg/ml 4',6'-diamino-3-phenylindole (DAPI) after washing with 50 mM sodium phosphate buffer (pH 7.0). Histology of seedlings was examined after fixation with FAA solution, as described previously by Tsukaya et al. (1993) and embedding in Technovit 7100 resin (Kulzer & Co. GmbH, Wehrheim, FRG).

tracted plasma. *J.*

jection reflex. *Curr.*

'olecular Cloning: A  
or, New York: Cold

sion of multiple fork  
rn formation in the

nization of forebrain  
of the rat. *J. Comp.*

nnell, S. M., Arias,  
l Rosenfeld, M. G.  
hypothalamus and  
lomain factor *Brn2*.

it for *Lim1* in head-

ll, A., Gardner, H.,  
nberg, R. A. (1995).  
oncogenesis in the

E., Norman, D. J.,  
A. K., Miller, A. P.,  
and Rosenfeld, M.  
: *Prophet of Pit-1*  
re 384, 327-333.

halamic integration:  
nuclei. *Ann. Rev.*

Bronson, R. T. and  
penia in mice with a  
ll 65, 1153-1163.  
y development: short

ess, P., Preziosos, V.  
ion factor *HNF-3β* is  
o. *Cell* 78, 575-588.  
an B. L. M. (1995).  
lorm formation and

, V. C. and Lai, E.  
essential for the  
41-1152.

mic connections: An  
. 201-215.

mouse *Bmp8a* (*Op2*)  
permatogenesis and

24 January 1997)

Incorporation of 5-bromodeoxyuridine (BrdU) into DNA was monitored after floating seedlings on a solution of 1 mM BrdU and 100  $\mu$ M 5-fluoro-5'-deoxyuridine for 13 hours at 23°C. Fixation and embedding of samples in Technovit 7100 resin were performed as described by Fujie et al. (1994). The samples, embedded in resin, were sectioned at 1.2  $\mu$ m thickness with glass knives. For detection of cells that had incorporated BrdU into their DNA, samples were treated with a BrdU-specific mouse monoclonal antibody (Becton Dickinson Immunochemistry Systems, San Jose, CA, USA) and sheep antibodies against mouse immunoglobulin, conjugated with fluorescein isothiocyanate (Amersham Japan Co. Ltd., Tokyo, Japan) as described elsewhere (Fujie et al., 1994).

For surgical experiments, one of the cotyledons was cut off with a razor blade at the base, at the stage indicated, and cultivation was continued. As a control to determine whether the death of seedlings could be due to damage resulting from the surgical treatment, whole seedlings were also grown on plates under sterile culture conditions.

For culture of segments of leaves in vitro, explants were cut into segments of approx. 1 cm in length and 1 cm in width with a razor blade. Explants were then put on MS101 medium (pH 6.3; Tsukaya et al., 1991) which was composed of Murashige-Skoog's inorganic salts supplemented with 1 mg/l benzyladenine as a cytokinin, 0.1 mg/l  $\alpha$ -naphthyl acetic acid as an auxin, 2% (w/v) sucrose, 3 mg/l thiamine-HCl, 5 mg/l nicotinic acid, 0.5 mg/l pyridoxine HCl, and 0.2 % Gellan gum (Wako Pure Chemical Industries, Ltd., Osaka, Japan). Cultiva-


tion was carried out at 23°C under continuous white light (at approx. 65  $\mu$ E/m<sup>2</sup>/s) as described by Tsukaya et al. (1994).


## RESULTS

The germination of *Monophyllaea horsfieldii* R. Br. was analyzed first. Embryos of *Monophyllaea* in mature seeds contained two cotyledons of equal size without a shoot apical meristem (Fig. 1A). Germinated seedlings also had two cotyledons of equal size (Fig. 1B,C). Seedlings lacked a shoot apical meristem (Fig. 1C). However, a meristem-like cluster of small cells, including mitotic cells, was located at the base of each half-blade of each cotyledon soon after germination (Fig. 1B). Since clusters of small cells are not necessarily indicative of mitotic tissues in leaf primordia (Dubuc-Lebreux and Sattler, 1980), the meristematic activity in the basal regions of the cotyledons was examined by monitoring the incorporation of BrdU into DNA. Evidence of DNA synthesis was detected only at the base of each cotyledon (Fig. 1D). By contrast, no cells at the S phase of the cell cycle were detected at the shoot apex (Fig. 1E).

Next, the differential growth of the two cotyledons was analyzed during plant development. After germination, the meristematic regions disappeared from one of the cotyledons

**Fig. 1.** Germination of seedlings of *Monophyllaea*. (A) A mature embryo of *Monophyllaea horsfieldii*. Bar, 100  $\mu$ m. (B) A young seedling just after germination. Note the absence of meristematic cells in the apical region of the embryo (indicated by an arrowhead). Arrows indicate meristematic regions at the base of the two cotyledons. Nuclei were stained with DAPI. Bar, 100  $\mu$ m. (C) Longitudinal section of a young seedling just after germination (see B). An arrowhead indicates the apical region, with no evidence of meristematic cells. Bar: 100  $\mu$ m. (D,E) Detection of DNA-synthesizing cells that had incorporated BrdU, at the base of the two cotyledons of a seedling just after germination. Nuclei emitting green fluorescent signals indicate cells that had incorporated BrdU at the S phase of the cell cycle. Arrows indicate nuclei with positive signals. Bar, 100  $\mu$ m. (D) Longitudinal section through the base of the two cotyledons. The upper photograph is a Nomarski image and the lower photograph was observed by a fluorescence microscopy. A vertical bar indicates the position of the shoot axis of the seedling (hypocotyl). The diagram shows the plane of the section in a seedling. A vertical dotted bar in the diagram indicates the position of the hypocotyl of the seedling. (E) Longitudinal section at the center of the seedling. The left photograph is a Nomarski image and right photograph is a fluorescence micrograph. (F) An adult specimen of *Monophyllaea horsfieldii*, with a single large leaf with inflorescences at its base. The photograph was taken in the plant's native habitat near Ipoh, Malaysia (Dec. 1995).






**Fig. 2.** Anisocotyledonous growth of *Monophyllaea* seedlings. (A) Variations in anisocotyledonous growth. Seedlings that had been cultivated on an agar plate were photographed 39 days after sowing. Bar, 5 mm. (B) Magnified view of the apical region of a seedling 15 days after sowing, showing the unequal distribution of meristematic cells in the two cotyledons. Note the disappearance of mitotic cells from the base of the left cotyledon. Bar, 100  $\mu$ m. (C) Lengths of the cotyledons were measured, and the length of the smaller one was plotted against that of the larger one for each seedling. Note the equal sizes of the cotyledons soon after germination (13 days after sowing, closed circles). The relationship between the length of smaller cotyledon (x) and that of larger cotyledon (y), at this stage, is represented by the equation shown in the figure, where r is the correlation coefficient. Note also the varied anisocotyledonous growth. Ages of seedlings examined (das, days after sowing) are indicated by different symbols as shown in the Figure.

(Fig. 2B) and anisocotyledonous development subsequently became apparent (Fig. 2A,C). The pattern of anisocotyledonous development varied significantly among individual plants (Fig. 2A,C), as mentioned previously by Oehlkers (1923), suggesting that such development might not be strictly determined in an 'all or none' manner. A small number of plants even had cotyledons of equal size 109 days after sowing (Fig. 2A,C).

How is the unequal development of the two cotyledons determined? Ninety years ago,

**Fig. 3.** Growth of cotyledons after surgical treatment. (A-D) Seedlings of *Monophyllaea* that had been cultivated on agar plates for 45 days. (A) A control plant. (B) A seedling from which one of the cotyledons had been removed 14 days after sowing, when both cotyledons were of equal size. (C) A seedling from which the smaller cotyledon had been removed 24 days after sowing. (D) A seedling from which the larger cotyledon had been removed 24 days after sowing. Bar, 5 mm. (E,F) Magnified views of seedlings that produced secondary cotyledons after removal of the larger cotyledon. (G) Scanning electron micrograph of primordia of secondary cotyledons that appeared at the base of the remaining cotyledon after surgical treatment. Arrowheads show primordia of secondary cotyledons. Bar, 100  $\mu$ m.




Chifflet (1909) noted that seedlings of *M. horsfieldii* occasionally developed a 'secondary cotyledon' after the larger cotyledon had been removed. In the present study, a re-examination of this phenomenon was made by surgical experiments at specific stages of development. When one of the two cotyledons was cut off soon after germination but before anisocotyledonous growth had become apparent, all the remaining cotyledons continued to develop (Table 1; Fig. 3B). This result implies that both cotyledons had the potential to grow continuously at a stage soon after germination. However, when one of the cotyledons was removed after a difference had become apparent in the sizes of the two cotyledons, different results were obtained. Seedlings from which smaller cotyledons had been removed continued to grow (Table 1; Fig. 3C). However, seedlings from which larger cotyledons had been removed retained a small cotyledon in many cases (Table 1; Fig. 3D). This result indicated that growth of smaller cotyledons had already been irreversibly interrupted. Oehlkers (1923) reported that some small cotyledons of seedlings, from which larger cotyledons had been surgically removed, continued to grow. However, in the present study, the percentage of determined cotyledons increased gradually with time after sowing (Table 1). Thus, the fate of the cotyledons, namely, the anisocotyledonous development of the seedling, seemed to be determined gradually after germination. It is noteworthy that, in many cases, a new leaf primordium differentiated irregularly at the base of the remaining cotyledon and/or on the hypocotyl (Table 2; Fig. 3D-G). This phenomenon was more frequently observed after the larger cotyledon had been removed than after the smaller one had been removed (Table 2). A similar phenomenon, with differentiation of secondary cotyledons, was also observed in a few seedlings without any surgical intervention (2 of 50 seedlings, Fig. 4).

To examine the role of the gravity in the determination of anisocotilyly, some seeds were sown on vertically oriented agar plates (Okada and Shimura 1992). However, the direction of the force of gravity seemed to have only a weak influence on

light (at approx.

R. Br. was mature seeds a shoot apical had two coty- a shoot apical cluster of small use of the each ion (Fig. 1B). indicative of ix and Sattler, s of the cotyle- on of BrdU into nly at the base at the S phase (Fig. 1E). cotyledons was mination, the he cotyledons



**Table 1A. The fate of the remaining cotyledons of seedlings that were grown on rockwool for 30 days after surgical treatment**

Day	% of remaining cotyledons that continued to grow		
	Removal of 1 of 2*	Removal of larger	Removal of smaller
13	100.0 (24)	—	—
15	61.1 (18)	—	—
16	46.5 (43)	—	—
20	—	0.0 (13)	100 (13)
24	—	17.6 (17)	100 (8)
27	—	6.7 (15)	100 (10)

\*Cotyledons were of equal size from day 13 to day 16.

Numbers in parentheses are the number of seedlings examined in each case.

**Table 1B. The fate of the remaining cotyledons of seedlings that were grown on plates for 39 days after surgical treatment**

Day	% of remaining cotyledons that continued to grow		
	Removal of 1 of 2*	Removal of larger	Removal of smaller
14	96.6 (29)	—	—
19	—	25.0 (20; 4 dead)	65.0 (20; 1 dead)
24	—	8.0 (25)	93.9 (33)

\*Cotyledons were of equal size on day 14.

Numbers in parentheses are the numbers of seedlings examined in each case.

**Table 2. The development of secondary cotyledons after surgical treatment of seedlings grown on plates for 39 days**

Day	% plants with secondary cotyledons following surgery		
	Removal of 1 of 2*	Removal of larger	Removal of smaller
14	13.8 (29)	—	—
19	—	5.0 (20; 4 dead)	5.0 (20; 1 dead)
24	—	72.0 (25)	3.0 (33)

\*Cotyledons were of equal size at 14 days.


Numbers in parentheses are the numbers of seedlings examined in each case.

the gravitropism of hypocotyls, without any affect on the direction of the anisocotyledonous growth (Fig. 5).


As a first step towards understanding of the roles of phytohormones in the regulation of leaf morphogenesis in *Monophyllaea*, segments of leaves were cultivated in vitro on medium supplemented with 1 mg/l. benzyladenine as a cytokinin and 0.1 mg/l  $\alpha$ -naphthyl acetic acid as an auxin. These conditions stimulate regeneration of shoots from explants of leaves of *Arabidopsis* (Tsukaya et al., 1991). As shown in Fig. 6A, explants differentiated many leaf-like structures, which resembled secondary cotyledons, on the surface of the explants. Scanning electron microscopy indicated that single secondary cotyledons seemed to occur independently of each other, without any shoot-like structures (Fig. 6B).

## DISCUSSION

Apical dominance is the term used for the control exerted by the apex of the shoot over the outgrowth of lateral buds and it is observed in most higher plants (Cline, 1991). Apical



**Fig. 4. Spontaneous occurrence of secondary cotyledons. At the base of the larger cotyledon, secondary cotyledons (one of them is indicated by an arrow) occurred spontaneously. Two of 50 seedlings exhibited this phenomenon 4 months after sowing, without any surgical treatment. Bar, 5 mm.**



**Fig. 5. Relationship between anisocotylly and the direction of the force of gravity. (A) Seedlings, 34 days after sowing that had been cultivated on a vertically oriented agar plate. The direction of the force of gravity is indicated by an arrow. (B) Summary of the directions in which the larger cotyledon elongated. The direction of the force of gravity is indicated by an arrow.**

dominance is maintained in a plant shoot only when the apical meristem of the shoot actively suppresses the growth of lateral meristem. When the apical meristem is damaged, competition occurs among the remaining lateral meristems, with the winner again establishing apical dominance. In such plants, competition and suppression control the development of shoots. From the present study, it appears that the phenomenon of anisocotyledonous development of the genus *Monophyllaea* is analogous to apical dominance, as will be discussed below.

In this study, several previously reported observations on one-leaf plants were confirmed in *Monophyllaea horsfieldii*, namely, the absence of a shoot apical meristem, the phenomenon of anisocotyledonous growth, and the presence of meristematic regions in the larger cotyledon. The locations of meristematic regions were demonstrated for the first time by direct monitoring of DNA synthesis. The following new findings were also made in this study. (1) Just after germination, the two cotyledons are completely identical morphologically. (2) Both cotyledons



ons. At the base  
them is  
of 50 seedlings  
ithout any




Fig. 6. Regeneration of secondary cotyledons on leaf segments in culture in vitro. (A) Occurrence of secondary cotyledons on the surface of a leaf segment. An explant after cultivation for two months on a plate of solidified medium is shown (see also Materials and Methods). (B) Scanning electron micrograph of a secondary cotyledon induced during culture in vitro. Note the single leaf-like organ without any structure, which resembles a shoot apical meristem at the base. Bar, 100  $\mu$ m.

have meristematic regions, at the base of the blade, just after the germination. This conclusion strongly supports the hypothesis proposed by Oehlkers (1923) that the two cotyledons are physiologically equivalent. Oehlkers (1923) observed varied patterns of anisocotylly and performed some surgical experiments. Jong and Burt (1975) suggested that, in *Streptocarpus fanniniae*, the meristematic region at the base of the larger cotyledon is the former embryonic meristem that has moved from its original position after the rapid growth of the larger cotyledon. However, this interpretation is not valid in the genus *Monophyllaea*. No embryonic meristem was recognized in mature embryos of *Monophyllaea*, and the meristematic regions at the base of the lamina were found only just after germination. Since anisocotyledonous growth is not necessarily accompanied by the absence of a shoot apical meristem in Gesneriaceae (as it is, for example, in species in the genera *Didymocarpus*, *Saint-paulia* and *Oreocharis*; Hill, 1938), absence of a shoot apical meristem can be considered to be independent of anisocotylly. (3) Anisocotyledonous growth is determined during growth of cotyledons and is irreversible. Prior to this study, the timing and mechanism of anisocotylly were unknown. Oehlkers (1923) suggested that the smaller cotyledons at later stages might still have ability to grow continuously. However, the present study strongly supports the irreversible determination of the fate of cotyledons at somewhat earlier stages. The difference between the conclusions might be due to the fact that Oehlkers succeeded in surgical experiments with only limited numbers of seedlings because of the high death rate of seedlings after surgical




Fig. 7. Model of the mechanism responsible for the developmental control of anisocotylly in *Monophyllaea*. Areas with dense dots represent meristematic regions. Arrows on leaf blades indicate further growth. See text for details.

treatment. Oehlkers did not present quantitative results of his surgical experiments. In the present study, the combination of sterile culture with surgical treatment resulted a very low death ratio of treated seedlings. Thus, the surgical experiments in the present study showed for the first time that both cotyledons have equal potential for continuous growth just after germination. The present study proved for the first time that anisocotyledonous growth is irreversible. Moreover, the present study also showed for the first time that the timing of the choice of dominant cotyledon is not strictly programmed.

The surgical experiments suggested that the two cotyledons compete with one another. The determination of anisocotyledonous development seemed to be the result of competition between the two cotyledons for the following reasons: (1) Surgical detachment of one cotyledon at an early stage resulted in further growth of all of the remaining cotyledons. (2) Determination of anisocotyledonous growth is not strictly programmed (in some plants both cotyledons developed equally). Fig. 7 shows a proposed model of the developmental control of anisocotylly in *Monophyllaea*, as deduced from this study. Just after germination, both cotyledons are identical and each has the ability to grow continuously. Cotyledons compete with one another and one finally inhibits the further growth of the other, perhaps winning the competition as a result of the influence of some environmental factor. The model shown in Fig. 7, which suggests that one cotyledon inhibits the growth of the other cotyledon, is supported by the fact that seedlings with equal-sized cotyledons tended to grow more slowly than seedlings with typical anisocotyledonous development (data not shown). There remains the possibility that 'self-activation' of cotyledons might play a role in the regulation of anisocotyledonous growth. With respect to the roles of phytohormones in anisocotylly, Rosenblum and Basile (1984) studied the effects of phytohormones on anisocotyledonous growth of some species in the genus *Streptocarpus*. They showed that some hormones, when supplied exogenously, influenced the fate of the cotyledons and the apical meristems in *Streptocarpus*. Since the combination of a cytokinin and an auxin resulted in differentiation of secondary cotyledons on the surface of leaf segments (Fig. 6), phytohormones might be involved in the regulation of anisocotylly. However, the roles of endogenous

phytohormones in anisocotylly cannot be discussed on the basis of experiments with exogenously supplied phytohormones.

The present study showed that the direction of the force of gravity had no effect on the determination of anisocotyledonous development. Oehlkers (1923) suggested the possibility that growth of cotyledons above the horizontal line might be promoted. However, in the present study, no results clearly supported this hypothesis. Further analysis is needed to define the physiological nature of the competition between the two cotyledons in *Monophyllaea*.

'Competitive organogenesis' has not been reported in other organisms, but this phenomenon might be analogous to the apical dominance that is commonly observed during the regulation of development of plant shoot systems. When one of the cotyledons was cut off, a secondary cotyledon frequently appeared at the base of the remaining cotyledon and/or on the hypocotyl, as noted by Chiffot (1909). Although this phenomenon might be a secondary effect induced by wounding stress, the appearance of a secondary cotyledon from the remaining cotyledon and/or hypocotyl can also be explained by the model shown in Fig. 7 (right). When the two cotyledons inhibit the meristematic activity of the other part of shoot and then the inhibitory effect of one of the cotyledons is abruptly terminated, the meristematic activity of the remaining cotyledon and the potential meristems in the hypocotyl might be enhanced. The appearance of secondary cotyledons might be the result of such sudden release from inhibition. The model is supported by the tendency of secondary cotyledons to appear more frequently when the larger cotyledon had been detached than when the smaller cotyledon had been detached (Table 2; Fig. 3D). This explanation is also applicable to the spontaneous differentiation of secondary cotyledons at the base of larger cotyledons (Fig. 4).

The present results suggests that, although the one-leaf phenotype of *Monophyllaea* is unusual, the mechanism that governs anisocotylly might be based on a principal that is common to and typical of all plant shoots, namely, competition and suppression. Apical dominance, which governs shoot morphogenesis in plants, is a more common example of competition and suppression. The plant-specific developmental principal that is based on competition and suppression can be considered to represent 'competitive organogenesis'. Thus, the anisocotyledonous phenotype is an extreme example of competitive organogenesis. Future studies of *Monophyllaea* should provide new insights into plant morphogenesis. The apparent similarity between seedlings of *Monophyllaea* and the seedlings of mutants of *Arabidopsis thaliana* that lack a shoot apical meristem (Barton and Poethig, 1993; Long et al., 1996; Laux et al., 1996) is also of interest in terms of the molecular mechanisms required for the establishment and maintenance of meristematic regions in plants.

The author thanks Dr M. Hayashi (The University of Tokyo) for guidance in the examination of the incorporation of BrdU. Prof. T. Kuroiwa (The University of Tokyo) kindly allowed the author to use an ultramicrotome in his laboratory. The author thanks the following people for critical discussions and/or help during studies carried out in the field and in the laboratory: Dr A.A. Bidin (University of Kebangsaan, Malaysia); Prof. M. Kato (The University of Tokyo); Dr H. Okada (Osaka University); Dr H. Setoguchi (Tokyo Metropolitan University); Mr M. Nakaya (The University of Tokyo); Prof. H. Uchimiya (The University of Tokyo); and Dr T. Tsuge (The Univer-

sity of Tokyo). The author also thanks Prof. K. Iwatsuki, the curator of the Botanical Garden, The University of Tokyo, for use of a scanning electron microscope. Prof. D. Okumoto (Saitama University) and Prof. A. Okada (Toyo University) helped the author to understand papers written in French and German. This study was supported by a Grant-in-Aid from the Monbusho International Scientific Research Program of the Ministry of Education, Science and Culture, Japan. The report by Oehlkers (1923) was brought to the author's attention by an anonymous referee.

## REFERENCES

- Barton, M.K. and Poethig, R.S. (1993). Formation of the shoot apical meristem in *Arabidopsis thaliana*: an analysis of development in the wild type and in the shoot meristemless mutant. *Development* **119**, 823-831.
- Bell, A. D. (1991) *Plant Form*. Oxford University Press, Oxford.
- Brown, R. (1840). *Monophyllaea*. In: Bennett, *Pl. Jav. Rar.* **121**.
- Burt, B. L. (1978). Studies in the Gesneriaceae of the Old World XLV: a preliminary revision of *Monophyllaea*. *Notes from the Royal Botanic Garden, Edinburgh* **37**, 1-59.
- Chiffot, M. (1909). Sur quelques variations du *Monophyllaea horsfieldii* R. Br. *Compt. Rend. Acad. Sci. Paris* **148**, 939-941.
- Cline, M. G. (1991). Apical dominance. *Bot. Rev.* **57**, 318-358.
- Dubuc-Lebreux, M. A. and Sattler, R. (1980). Développement des organes foliacés chez *Nicotiana tabacum* et le problème des méristèmes marginaux. *Phytomorph.* **30**, 17-32.
- Fujie, M., Kuroiwa, H., Kawano, S., Mutoh, S. and Kuroiwa, K. (1994). Behavior of organelles and their nucleoids in the shoot apical meristem during leaf development in *Arabidopsis thaliana* L. *Planta* **194**, 395-405.
- Hill, A. W. (1938). The monocotyledonous seedlings of certain dicotyledons. With special reference to the Gesneriaceae. *Ann. Bot. N. S.* **2**, 127-143.
- Jong, K. and Burt, B. L. (1975). The evolution of morphological novelty exemplified in the growth patterns of some Gesneriaceae. *New Phytol.* **75**, 297-311.
- Laux, T., Mayer, K. F., Berger, J. and Jurgens, G. (1996). The *WUSCHEL* gene is required for shoot and floral meristem integrity in *Arabidopsis*. *Development* **122**, 87-96.
- Long, J. A., Moan, E. I., Medford, J. I. and Barton, M. K. (1996). A member of the *KNOTTED1* class of homeodomain proteins encoded by the *STM* genes of *Arabidopsis*. *Nature* **379**, 66-69.
- Oehlkers, F. (1923). Entwicklungsgeschichte von *Monophyllaea horsfieldii*. *Beih. z. Bot. Centralbl. I. Abt.* **39**, 128-151.
- Okada, K. and Shimura, Y. (1992). Mutational analysis of root gravitropism and phototropism of *Arabidopsis thaliana* seedlings. *Aust. Plant Physiol.* **19**, 439-448.
- Rosenblum, I. M. and Basile, D. V. (1984). Hormonal regulation of morphogenesis in *Streptocarpus* and its relevance to evolutionary history to the Gesneriaceae. *Amer. J. Bot.* **71**, 52-64.
- Tsuge, T., Tsukaya, H. and Uchimiya, H. (1996). Two independent and polarized processes of cell elongation regulate leaf blade expansion in *Arabidopsis thaliana* (L.) Heynh. *Development* **122**, 1589-1600.
- Tsukaya, H., Naito, S., Rédei, G. P. and Komeda, Y. (1993). A new class of mutations in *Arabidopsis thaliana*, *acaulis1*, affecting the development of both inflorescences and leaves. *Development* **118**, 751-764.
- Tsukaya, H., Ohshima, T., Naito, S., Chino, M. and Komeda, Y. (1991). Sugar-dependent expression of the *CHS-A* gene for chalcone synthase from petunia in transgenic *Arabidopsis*. *Plant Physiol.* **97**, 1414-1421.
- Tsukaya, H., Tsuge, T. and Uchimiya, H. (1994). The cotyledon: a superior system for studies of leaf development. *Planta* **195**, 309-312.
- Wardlaw, C. W. (1952). *Phylogeny and Morphogenesis*. Macmillan and Co., Ltd, London.
- Weber, A. (1975). Beiträge zur Morphologie und Systematik der Klugieae und Loxonieae (Gesneriaceae). I. Die Sproß- und Infloreszenzorganisation von *Monophyllaea* R. Br. *Bot. Jahrb. Syst.* **95**, 174-207.
- Weber, A. (1976a) Beiträge zur Morphologie und Systematik der Klugieae und Loxonieae (Gesneriaceae). II. Morphologie, Anatomie und Ontogenese der Blüte von *Monophyllaea* R. Br. *Bot. Jahrb. Syst.* **95**, 435-454.
- Weber, A. (1976b) Beiträge zur Morphologie und Systematik der Klugieae und Loxonieae (Gesneriaceae). III. *Whytockia* als morphologische und phylogenetische Ausgangsform von *Monophyllaea*. *Beitr. Biol. Pflanzen* **52**, 183-205

(Accepted 30 January 1997)

PLEASE PRINT OR TYPE CLEARLY

ONE REQUEST PER FORM

Today's Date: 15 Jan. # 2002 Not Needed After (Date):

REQUESTOR'S NAME: Boggan, John

Library Card Number: 2-9088-\_\_\_\_\_

Smithsonian Mailing Address (please include MRC): Dept - Biology / Botany  
MRC 166

Phone Number: (202) 357-2306 E-Mail Address: boggan.john@si.edu

INSTRUCTIONS FOR FILLING OUT FORM

- fill out left side of form completely
- look up item on SIRIS ([www.siris.si.edu](http://www.siris.si.edu))
- write down call number and indicate all locations
- indicate "not on SIRIS" under Notes and Comments if you can't locate on SIRIS
- indicate "not on shelf" (if applicable)
- indicate circulation status (if known)

SIRIS Call No. QL 951 J865 IN SC. STRI

ITEM REQUESTED

Book or article author: Tsukaya, H.

Book title or journal name: ~~Determinator~~ Development

Article title, if applicable: Determination of the unequal fate of  
the copolymers in the one-leaved plant Mangrove

Pages: 1275-1280

Holding Libraries: (If item is in SIRIS) Circle/fill in all locations:

ANTH	ENT	MIN	NMAA/NPG	NZP
BIRDS	FISH	MOLL	NMAH	REPT
BOTANY	INV ZOO	MSC	NMAI	SERC
CRLS	MAMM	NASM	NMNH	SI-ANNEX
				STRI

OTHER LOCATIONS:

Notes and Comments:

OCLC Number (if known):

John -  
your other  
request (from  
Advances in genetics)  
could not be filled  
at MSC, as they lack  
the volume. I have  
given the request  
back to ILL to  
get from STRI  
or another source  
if they prefer  
Ruth

**NO COPYRIGHT RESTRICTIONS** The copyright law of the United States (Title 17, United States Code) governs the  
copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other  
the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a  
request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement. This institution reserves the right to refuse to  
accept a copying order if, in its judgement, fulfillment of the order would involve violation of copyright law (37 C.F.R. Paragraph 201.14).