

Luegmayr 1993a

Pollen of Hawaiian *Cyrtandra* (Gesneriaceae) including notes on Southeast Asian taxa.

Blumea 38: 25-38, fig. 1-5.


REFNO: 2186

KEYWORDS:

Borneo, *Cyrtandra*, Hawaii, Hybrid, Pollen

S
B
P

BLUMEA



Volume 38, No. 1
1993

JOURNAL OF PLANT TAXONOMY AND PLANT GEOGRAPHY

POLLEN OF HAWAIIAN CYRTANDRA (GESNERIACEAE)
INCLUDING NOTES ON SOUTHEAST ASIAN TAXA

EVA LUEGMAYR

Institute of Botany, University of Vienna, Rennweg 14, 1030 Vienna, Austria

SUMMARY

Pollen of 30 species and 12 hybrids (88 samples in total) of Hawaiian *Cyrtandra*, covering all 6 sections recognized on the Hawaiian archipelago, was examined using light, scanning and transmission electron microscopy. Hawaiian *Cyrtandra* is stenopalynous. The pollen grains are binucleate, isopolar, small-sized and 3-colporoidate (very rarely 4-colporoidate). In equatorial view they are spheroidal, in polar view circular or semiangular. The tectum is mostly microreticulate with brochi of different sizes (heterobrochate), very rarely reticulate or perforate. Supratectal elements are lacking. Some variation can be observed in number and diameter of the lumina. Palynological features of most of the investigated taxa tend to overlap and thus are not very helpful in supporting the infrageneric division of *Cyrtandra* on Hawaii.

Pollen viability of hybrids, which is said to be reduced occasionally, was examined and found to be high with rates of 70 to 99%. Therefore, reduced pollen viability is not a reliable character for designating hybrid status.

Asian *Cyrtandra* is eurypalynous. For this study, 19 species, originating from the Malay Peninsula and Borneo, have been investigated. Two exine types, differing in sculpture and structure, can be distinguished: 1) tectum microreticulate-reticulate, supratectal elements lacking, and 2) tectum microreticulate-perforate with numerous conical supratectal elements. The number, diameter and shape of lumina provide further important diagnostic features.

INTRODUCTION

Cyrtandra, the largest genus of the family Gesneriaceae, comprises about 500-600 species ranging from SE Asia throughout Malesia, NE Australia and the Pacific islands. Centres of diversity include New Guinea, Borneo, each with over 150 species, and the Philippines. The only treatment covering the whole genus dates back to Clarke (1883). However, Clarke's subdivision of the genus into sections is more or less ignored in nearly all subsequent regional treatments because of the confusing delimitation of the sections.

On the Hawaiian archipelago 53 species are recognized currently (Wagner et al., 1990). This means a drastic reduction of species number compared with former treatments, especially that of St. John (1966), who recognized and described 131 species only for the island of Oahu. The considerable discrepancy is due to a different taxonomic philosophy: while St. John delimited species very narrowly, Wagner et al. accepted a number of species to be very polymorphic. Moreover, Wagner et al. interpreted individuals exhibiting intermediate characters between two sympatric species as hybrids. To date, 67 putative hybrid combinations have been recorded (Wagner & Herbst, in prep.).

Table 1. List of investigated taxa of Hawaiian *Cyrtandra*, origins and short characteristics of the exine pattern (number of lumina per 25 μm^2 of the mesocolpia / diameter of the largest lumina of the mesocolpia in μm).

Taxa	Origin, voucher	Lumina number / size
Sect. <i>Crotonocalyces</i> Hillebr.		
<i>C. cordifolia</i> Gaud.	Oahu, Kiehn et al. 900722-5/1 (WU, PTBG) -, W. C. Gagne 523 (BISH)	120 / 0.5 80 / 0.8
<i>C. kealiae</i> Wawra	Kauai, St. John et al. 23076 (BISH) -, Kiehn & Luegmayer 920825-1/1 (WU)	90 / 0.5 120 / 0.7
<i>C. limahuliensis</i> St. John	Kauai, Kiehn et al. 900823-2/12 (WU, US)	120 / 0.8
<i>C. platyphylla</i> A. Gray	Hawaii, Kiehn & Cuddihy 900812-2/1 (WU) s.l., Warshauer 1933 (BISH) Hawaii, Warshauer 1275 (BISH) s.l., Warshauer 1585 (BISH)	100 / 0.6 50 / 0.4 70 / 0.6
<i>C. propinqua</i> C. Forbes	Oahu, Takeuchi 2955 (BISH) -, Takeuchi 3045 (BISH) -, Kiehn et al. 920909-3/2 (WU, US, BISH)	110 / 0.4 120 / 0.5
<i>C. wawrae</i> C. B. Clarke	Kauai, Wagner et al. 6064 (BISH)	50 / 0.9
Sect. <i>Apertae</i> C. B. Clarke		
<i>C. garnotiana</i> Gaud.	Oahu, Takeuchi 2142 (BISH) s.l., Obata 86-650 (BISH)	90 / 0.7 110 / 0.4
<i>C. laxiflora</i> H. Mann	Oahu, Kiehn & Obata 890809-1/6 (WU, BISH) -, Takeuchi 1779 (BISH) -, Takeuchi 1868 (BISH) -, Kiehn et al. 920909-3/1 (WU, US) s.l., Takeuchi 2532 (BISH)	180 / 0.2 100 / 0.3 60 / 0.7 200 / 0.2 140 / 0.6
<i>C. sandwicensis</i> (H. Lév.) St. John & Storey	Oahu, Kiehn et al. 900722-2/1 (WU, PTBG, BISH) -, Kiehn et al. 900722-2/3 (WU, PTBG, BISH) -, Kiehn 920907-1/9 (WU, US)	120 / 0.4 80 / 0.4 70 / 0.8
Sect. <i>Macrosepala</i> C. B. Clarke		
<i>C. grayana</i> Hillebr.	Maui, Kiehn et al. 890723-2/2 (WU) Lanai, Hobdy s.n. (BISH) Maui, Nagata 3558 (BISH) -, Warshauer 3044 (BISH)	90 / 0.3 60 / 0.8 60 / 0.5 60 / 0.6
<i>C. grayi</i> C. B. Clarke	Molokai, Nagata 1141 (BISH) Maui, Warshauer 3068 (BISH) -, Warshauer 3106 (BISH)	50 / 0.7 100 / 0.4 110 / 0.4
<i>C. hashimotoi</i> Rock	Maui, Kiehn et al. 900803-3/3 (WU, US) -, Higashino 9319 (BISH) -, Hobdy 1894 (BISH)	50 / 0.6 40 / 0.3 80 / 0.7
<i>C. kauaiensis</i> Wawra	Kauai, Kiehn & Luegmayer 920825-3/3 (WU) -, Kiehn & Luegmayer 920830-4/1 (WU) s.l., Christensen 209 (BISH)	140 / 0.2 80 / 0.6
<i>C. lysiosepala</i> (A. Gray) C. B. Clarke	Hawaii, Kiehn & Cuddihy 900812-2/3 (WU, US) s.l., Cuddihy 216 (BISH) Hawaii, Warshauer 1653 (BISH) -, Warshauer 1668 (BISH) -, Warshauer 1656 (BISH)	80 / 0.7 150 / 0.3 120 / 0.6

(Table 1 continued)

Taxa	Origin, voucher	Lumina number / size
<i>(Sect. Cyllindrocalyces)</i>		
<i>C. macrocalyx</i> Hillebr.	Molokai, Kiehn & Luegmayer 920817-212 (WU)	80 / 0.6
<i>C. procera</i> Hillebr.	Molokai, Kiehn & Cuddihy 900812-213 (WU)	90 / 0.8
	s.l., Warshauer 2385 (BISH)	70 / 0.6
<i>C. cf. procera</i>	Molokai, Kiehn et al. 900806-312 (WU)	110 / 0.5
	-, Kiehn et al. 900806-211 (US)	90 / 0.4
<i>C. spathulata</i> St. John	Maui, Wagner et al. 5847 (BISH)	70 / 0.4
	-, Kiehn & Luegmayer 920821-211 (WU)	
<i>C. wainihaensis</i> H. Lév.	Kauai, Christensen 270 (BISH)	60 / 0.8
<i>Sect. Chaetocalyces</i> Hillebr.		
<i>C. giffardii</i> Rock	Hawaii, Warshauer 1647 (BISH)	40 / 0.2
<i>C. lydgatei</i> Hillebr.	Molokai, Nagata 1144 (BISH)	60 / 0.3
<i>C. macraei</i> A. Gray	Oahu, Nagata 3281 (BISH)	
<i>C. menziesii</i> Hook. & Arnott	Hawaii, Davis s.n. (BISH)	80 / 0.3
	-, Kiehn & Luegmayer 920821-111 (WU, PTBG)	40 / 0.6
<i>Sect. Verticillatae</i> St. John		
<i>C. calpidicarpa</i> (Rock)	Oahu, Takeuchi 2404 (BISH)	60 / 0.8
St. John & Storey	-, Kiehn et al. 920909-313 (WU, PTBG, BISH)	80 / 0.9
<i>C. confertiflora</i> (Wawra)	Kauai, Christensen 251 (BISH)	50 / 1.0
C.B. Clarke	-, Perlman 474 (BISH)	70 / 0.5
<i>C. grandiflora</i> Gaud.	Oahu, Kiehn et al. 900722-111 (WU, PTBG, BISH)	40 / 0.7
	-, Funk 141 (BISH)	50 / 1.3
	-, Kiehn 920907-112 (WU, US)	80 / 0.8
<i>C. hawaiiensis</i> C.B. Clarke	Oahu, Kiehn & Obata 890809-113 (WU)	70 / 0.9
	Maui, Wagner et al. 5855 (BISH)	60 / 0.9
	Hawaii, Kiehn & Luegmayer 920819-111 (WU)	80 / 0.8
	s.l., St. John 21360 (BISH)	40 / 1.1
<i>C. cf. hawaiiensis</i>	Oahu, Kiehn & Obata 890809-115 (WU)	80 / 0.8
<i>C. kaulantha</i> St. John & Storey	Oahu, Kiehn & Obata 890809-211 (WU)	50 / 0.7
<i>C. oenobarba</i> H. Mann	Kauai, Kiehn et al. 900823-113 (PTBG)	70 / 0.6
<i>Sect. Cyllindrocalyces</i> Hillebr.		
<i>C. longifolia</i> (Wawra)	Kauai, Kiehn & Flynn 900907-1113 (US)	70 / 0.5
Hillebr. ex C.B. Clarke		
<i>C. paludosa</i> Gaud.	Hawaii, St. John 23927 (BISH)	60 / 1.0
	Maui, Kiehn et al. 900803-711 (WU, US)	80 / 0.7
	Oahu, Takeuchi & Imada 1835 (BISH)	80 / 0.7
var. <i>paludosa</i>	s.l., Higashino 8015 (BISH)	60 / 0.9

Table 2. List of investigated hybrids of Hawaiian *Cyrtandra*, origins, short characteristics of the exine pattern (number of lumina per 25 μm^2 of the mesocolpia / diameter of the largest lumina of the mesocolpia in μm) and percentages of viable pollen grains. — Symbols for the sections: ● = *Crotonocalyces*, ○ = *Apertae*, ■ = *Macrosepalae*, □ = *Chaetocalyces*, ▼ = *Verticillatae*, ▽ = *Cylindrocalyces*.

Hybrid combinations	Origin, voucher	Lumina Number / Size	Pollen viability (%)
▼ <i>C. confertiflora</i> × <i>C. wainihaensis</i> ■	s.l., Perlman 475 (BISH)	50 / 0.7	89
● <i>C. cordifolia</i> × <i>C. sandwicensis</i> ○	Oahu, Kiehn 920907-1/18 (WU)	110 / 0.6	
□ <i>C. giffardii</i> × <i>C. platyphylla</i> ●	Hawaii, Nagata 2930 (BISH)	60 / 0.4	94
	—, St. John 22309 (BISH)	50 / 0.6	82
▼ <i>C. grandiflora</i> × <i>C. cordifolia</i> ●	Oahu, Kiehn 920907-1/2a (WU)	90 / 0.6	99
▼ <i>C. grandiflora</i> × <i>C. paludosa</i> ▽	Oahu, Takeuchi 3044 (BISH)	50 / 1.4	96
■ <i>C. grayi</i> × <i>C. platyphylla</i> ●	s.l., St. John et al. 25741 (BISH)	110 / 0.3	70
▼ <i>C. hawaiiensis</i> × <i>C. menziesii</i> □	Hawaii, Kiehn & Luegmayr 920819-1/2 (WU)	70 / 0.5	97
	s.l., Warshauer 1932 (BISH)	40 / 0.3	98
○ <i>C. laxiflora</i> × <i>C. paludosa</i> ▽	Oahu, Takeuchi 3049 (BISH)	90 / 0.4	92
▽ <i>C. paludosa</i> × <i>C. hashimotoi</i> ■	Maui, Kiehn et al. 900803-7/2 (WU)		
▽ <i>C. paludosa</i> × <i>C. platyphylla</i> ●	s.l., Davis 689 (BISH)		82
● <i>C. propinqua</i> × <i>C. calpidicarpa</i> ▼	Oahu, Kiehn et al. 920909-3/4 (WU)	60 / 0.7	2
	—, Kiehn et al. 920909-3/4b (WU)	60 / 0.7	96
	—, Takeuchi 2399 (BISH)	80 / 0.7	98
● <i>C. propinqua</i> × <i>C. garnotiana</i> ○	Oahu, Kiehn et al. 920909-4/1 (WU)	180 / 0.3	94

Apart from a single taxon (*C. grandiflora*, Roelofs, 1980) palynological information is lacking for Hawaiian *Cyrtandra*. The present study includes 30 species and 12 hybrids, using light, scanning and transmission electron microscopy. The investigation covers representatives of all six sections recognized on the Hawaiian archipelago (see Tables 1 & 2).

In order to evaluate the taxonomic significance of the data obtained and to provide additional palynological information on the genus, Hawaiian *Cyrtandra* is compared with SE Asian taxa. So far 19 species, originating from Borneo and the Malay Peninsula, have been studied (see Table 3); preliminary results of this investigation are provided.

MATERIAL AND METHODS

Pollen samples of Hawaiian taxa were taken either from alcohol- or glutaraldehyde-preserved flowers obtained from BISH and WU, or from plants grown in the greenhouses at the Botanical Garden of the University of Vienna (HBV). For the investigation of Asian taxa only herbarium material was available, except of *Cyrtandra pendula*, which is cultivated at HBV. Because of the marked protandry, flowers shortly

igins, short characteristics of the diameter of the largest lumina of : — Symbols for the sections: *haetocalyces*, \blacktriangledown = *Verticillatae*,

	Lumina Number / Size	Pollen viability (%)
H)	50 / 0.7	89
18 (WU)	110 / 0.6	
BISH)	60 / 0.4	94
3H)	50 / 0.6	82
12a (WU)	90 / 0.6	99
BISH)	50 / 1.4	96
741 (BISH)	110 / 0.3	70
<i>gmayr</i>		
	70 / 0.5	97
(BISH)	40 / 0.3	98
(BISH)	90 / 0.4	92
803-712 (WU)		
H)		82
20909-314		
	60 / 0.7	2
19-314b (WU)	60 / 0.7	96
SH)	80 / 0.7	98
)	180 / 0.3	94

s, 1980) palynological information includes 30 species and 12 ron microscopy. The investigation on the Hawaiian archipelago

the data obtained and to provide lawaiian *Cyrtandra* is compared m Borneo and the Malay Penin- results of this investigation are

ODS

from alcohol- or glutaraldehyde- from plants grown in the green-Vienna (HBV). For the investigatable, except of *Cyrtandra pen-* marked protandry, flowers shortly

Table 3. List of investigated taxa of SE Asian *Cyrtandra*, origins and short characteristics of the exine pattern (supratracteal elements present (+) or absent (-); number of lumina for 25 μm^2 of the mesocolpia / diameter of the largest lumina in μm and the presence (+) or absence (-) of rod-like luminal elements. Abbreviations: Sculpt.= supratracteal elements; * = lumina at the apocolpia markedly larger and fewer in number. Sectional allocation according to Clarke (1883) / suggested allocation to sections by Burt (pers. comm.). Burt replaced sectional names by ? wherever the species had been wrongly assigned.

Origin, Taxa	Sectional allocation according to		Lumina No./Size	Luminal Elements	Voucher
	Clarke	Burt			
Malay Peninsula					
<i>C. cupulata</i> Ridley	?	Disparae C.B. Clarke	50/1.3	+	W. & A. ¹ 870519-113 (WU)
<i>C. dispar</i> DC.		prob. <i>Decurrentes</i> C.B. Clarke	40/1.6	+	Weber 860818-212 (WU)
<i>C. pendula</i> Blume		<i>Decurrentes</i> C.B. Clarke	130/0.8	-	Weber 870501-1117 (WU)
<i>C. wallitchii</i> (C.B. Clarke) B.L. Burt			140/1.1	-	W. & A. 860828-114 (WU)
Borneo					
<i>C. farinosa</i> C.B. Clarke			100/0.7	-	Weber 790905-118 (WU)
<i>C. radiceflora</i> C.B. Clarke			70/0.8	-	Burt & Woods B1948 (E)
<i>C. basiflora</i> C.B. Clarke			*110/0.8	+	Argent et al. 753 (E)
<i>C. multibracteata</i> C.B. Clarke		<i>Dissimiles</i> C.B. Clarke		-	Burt & Martin B4801 (E)
<i>C. cretacea</i> Kraenzl.			110/0.9	-	Burt & Woods B1991 (E)
<i>C. chrysea</i> C.B. Clarke				-	Weber 790922 (WU)
<i>C. eximia</i> C.B. Clarke			80/0.8	-	Burt & Woods B1941 (E)
<i>C. oblongifolia</i> (Blume) C.B. Clarke		<i>Whittia</i> (Blume) C.B. Clarke	280/0.3	-	Argent et al. 742 (E)
<i>C. sarawakensis</i> C.B. Clarke			50/0.6	-	Burt & Martin B4712 (E)
<i>C. splendens</i> C.B. Clarke			*130/0.5	+	Burt & Martin B4815 (E)
<i>C. clarkei</i> Stapf			70/0.8	-	Sinclair 9104 (E)
			60/0.7	-	Weber 790924 (WU)
			60/0.9	-	Burt B8268 (E)
<i>C. gibbsiae</i> S. Moore			70/1.1	-	Burt & Martin B4945 (E)
<i>C. gillettiana</i> B.L. Burt			190/0.5	-	Burt & Martin B5169 (E)
<i>C. horizontalis</i> B.L. Burt		<i>Whittia</i> (Blume) C.B. Clarke		-	Burt & Woods B2579 (E)
<i>C. woodsii</i> B.L. Burt		<i>Decurrentes</i> C.B. Clarke	180/0.6	-	

¹⁾ Abbreviation for Weber & Anthonyasamy.

before or immediately after anthesis are in the best stage for pollen examination. Taxa and collections used for detailed investigations are listed in Table 1 (Hawaiian species), Table 2 (Hawaiian hybrids) and Table 3 (Asian species). The nomenclature follows Clarke (1883), Burt (pers. comm.) and Wagner et al. (1990).

For LM investigations fresh (living) pollen was mounted in glycerine jelly according to Wodehouse (1935). Pollen of hybrids was stained in acetocarmine for determining the nuclear condition and viability. Ratio of pollen viability is based on respective counts of 100 pollen grains of each sample.

For SEM investigations anthers of dried flowers were soaked in a 10% aqueous mixture of dioctyl sodium sulfosuccinate and 95% acetone for 24 hours (Peterson et al., 1978). Soaked and/or fixed anthers were first dehydrated in an alcohol series, subsequently chemically dehydrated in FDA (formaldehyde-dimethyl-acetal; Gerstenberger & Leins, 1978) and then critical-point-dried. A few samples were acetolyzed following the method of Erdtman (1960). The dried pollen grains were placed on aluminium specimen stubs and sputter-coated with gold.

For TEM investigations fixed anthers were dehydrated in an ethanol series and embedded in Spurr's mixture (Spurr, 1969). The ultrathin sections were stained with uranyl acetate and lead citrate. For detecting neutral polysaccharides ultrathin sections were treated with periodic acid-thiocarbohydrazide-silver protein (PA-TCH-SP) according to Thiéry (1967).

Width and amount of lumina per $25 \mu\text{m}^2$ were measured on SEM graphs. Pollen terminology follows Erdtman (1969), Faegri & Iversen (1975), and Pragłowski & Punt (1973).

RESULTS

Pollen description of Hawaiian taxa

The pollen grains, binucleate at maturity (Fig. 4A), are isopolar, 3-colporoidate (very rarely 4-colporoidate) and spheroidal (rarely suboblate) in equatorial view (Figs. 1B, 2D, 3C). In polar view the grains are circular or semiangular (Fig. 1D). Average equatorial diameters range from 14 to 23 μm .

The colpi are long and \pm tapering; the endexinous colpus membrane is covered with ectexinous granular elements (Figs. 2E, 4A). The colpus membrane is splitting up into lamellae at the equatorial part (Fig. 1E) and, after acetolysis, most colpi show an irregular os in this region (cf. Fritze & Williams, 1988; Luegmayr, 1993). However, endoapertures are not visible in the LM and there is no distinct endosculpturing at the aperture region. Thus, the aperture system is designated as colporoidate in the sense of Erdtman (1952).

The exine patterns at the mesocolpia are mostly microreticulate with minute muri and lumina less than 1 μm in width (Figs. 1C, 2C, 3B, 3D), very rarely reticulate or perforate (see Table 1). Lumina width varies considerably within the mesocolpia of a single grain (e.g. between 0.6 and 0.05 μm in *C. platyphylla*; Fig. 1C). Thus the microreticula are heterobrochate sensu Erdtman (1952). Moreover, the width of the lumina generally decreases towards the polar region and the colpus border (Figs. 1A and 3A show pollen grains with perforate apocolpia). Lumina irregular or circular, without free standing elements.

ge for pollen examination. Taxa listed in Table 1 (Hawaiian species), etc.). The nomenclature follows (1990).

mounted in glycerine jelly according to standard procedures. Pollen viability is based on re-

were soaked in a 10% aqueous acetone for 24 hours (Peterson 1988), dehydrated in an alcohol series, cleared in cedar oil, and mounted in glycerine jelly. A few samples were acetolyzed. Pollen grains were placed on slides.

stained in an ethanol series and thin sections were stained with uranyl acetate. Ultrathin sections were stained with lead citrate. Silver protein (PA-TCH-SP) was used for immunogold labelling.

Micrographs were prepared on SEM graphs. Pollen viability was determined (1975), and Pragłowski &

are isopolar, 3-colporoidate (oboblate) in equatorial view or circular or semiangular (Fig. 1D).

colpus membrane is covered with granular elements. The colpus membrane is splitting after acetolysis, most colpi show a distinct endosculpturing (Peterson 1988; Luegmayr, 1993). However, there is no distinct endosculpturing in taxa designated as colporoidate in the

microreticulate with minute muri (Fig. 3D), very rarely reticulate or microreticulate within the mesocolpia of a *typhylla*; Fig. 1C). Thus the aperture is very narrow. Moreover, the width of the colpi varies within the colpus border (Figs. 1A, 1B). Lumina irregular or circular,

Supratectal elements are lacking. A fine granular ornamentation, delicate cracks and shallow grooves radiating from the lumina are visible in Figures 1C, 3D and 3C. Except of the latter, these features are probably of artificial nature.

The exine consists of an ectexine (tectum, columellae and foot layer) and an endexine (Fig. 2A). The tectum is thicker than the foot layer; the columellate layer is at least as thick as the tectum thickness. The endexine is thin at the mesocolpia, but increases in thickness at the aperture region, forming a colpus membrane. The intine is 2-layered; the exintine is thickened at the aperture region with radially oriented and transverse channel-like structures (Fig. 1E).

Noteworthy is the occurrence of numerous pollenkitt droplets (Figs. 2B, 3E, 3F). Occasionally crystals cover the pollen grain surface and the inner wall of the anthers (Fig. 4C, E).

Hawaiian hybrids

No significant palynological differences have been found between 'good' species and hybrids, thus the former pollen description is valid for the examined hybrids also. Apart from a single individual of *C. propinqua* × *C. calpidicarpa*, pollen viability is high in hybrids with rates of 70 to 99% (see Table 2 & Fig. 4B, D). Abnormal, infertile pollen grains do not seem to occur more often in supposed hybrids.

One has to note that hybridisation primarily takes place between species of different sections, intrasectional hybrids are rare (see Table 2, cf. Wagner et al., 1990).

Brief palynological description of SE Asian taxa

Pollen grains 3-colporoidate (very rarely 4-colporoidate); spheroidal, outline in polar view circular to semiangular (Fig. 5A, C). Average equatorial diameters: 14–20 µm.

Colpi long and ± tapering; endexinous colpus membrane covered with ectexinous granular elements (Fig. 5B).

Exine pattern: reticulate (Fig. 5G), microreticulate (Fig. 5D, F) or almost perforate (Fig. 5E); supratectal elements present (Fig. 5A, B, D) or absent (Fig. 5C, E–G). Lumina occasionally with rod-shaped elements (Fig. 5G).

Exine consisting of an ectexine (tectum, columellae and footlayer) and an endexine. In *C. pendula* (cf. Luegmayr, 1993) the tectum is three times thicker than the footlayer. Columellate layer thinner than the tectum; supratectal elements conical, 0.2 µm high. Lamellated endexine thin at the mesocolpia, but increasing in thickness at the aperture region, forming a colpus membrane. Intine 2-layered.

DISCUSSION

Hawaiian taxa of *Cyrtandra* are stenopalynous. The pollen grains resemble one another strongly in size (14–23 µm), shape (spheroidal), apertures (3-colporoidate, very rarely 4-colporoidate) and exine ornamentation (microreticulate, very rarely reticulate or perforate; supratectal elements are lacking). Lumina width varies considerably within the mesocolpia of a single grain (e.g. between 0.8–0.08 µm in *C. procera*, Fig. 2C), thus the microreticula (or reticula) are heterobrochate sensu

Erdtman (1952). Moreover, the width of lumina generally decreases towards the polar region and the colpus border. Some variation also occurs in the number of tectal lumina (see Table 1 & 2). The number of lumina per $25 \mu\text{m}^2$ ranges from 40 (*C. giffardii*, *C. grandiflora*, *C. hashimotoi*, *C. menziesii*) to 200 (*C. laxiflora*). It is noticed that lumina number is not a constant character within a given species; intra-specific variation is up to 100% (cf. Table 1). Nevertheless, a slight trend can be noticed: pollen grains of taxa of the sections *Crotonocalyces*, *Apertae* and *Macrosepalae* tend to have more lumina (70–200 per $25 \mu\text{m}^2$ at average, rarely 40–60) than taxa of the sections *Chaetocalyces*, *Verticillatae* and *Cylindrocalyces* (40–80 per $25 \mu\text{m}^2$). In general, the palynological data obtained are not helpful in supporting the infrageneric delimitation of Hawaiian *Cyrtandra*.

In contrast to the uniform pollen grains of Hawaiian *Cyrtandra*, SE Asian *Cyrtandra* is eurypalynous. Pollen grains of 19 taxa, originating from the Malay Peninsula and Borneo show two different exine patterns, differing in sculpture and structure: 1) tectum microreticulate-reticulate, suprategal elements lacking, and 2) tectum microreticulate-perforate with numerous conical suprategal elements. The number, diameter and shape of lumina provide further important diagnostic features (see Table 3). The taxonomic value of the present data is not yet assessable since the infrageneric classification of the large variable genus is still in a poor state. A very first, interesting result is, that the few species that Burtt allocated to the sections *Whitia* and *Decurrentes* are palynologically homogeneous by having suprategal elements and a large number of lumina per $25 \mu\text{m}^2$ (cf. Table 3). Pollen morphology might be a helpful additional character to find out links with Hawaiian taxa but more data on Asian and Pacific taxa are needed to make more precise statements.

According to Wagner et al. (1990) hybridization is a widespread phenomenon in Hawaiian *Cyrtandra*. One criterion for designating hybrids is the occasionally reduced pollen fertility (actually the authors observed pollen viability by staining the pollen with Alexander's stain; cf. Alexander, 1969). Reduced pollen fertility, however, has not been constantly met, e.g. the hybrid between *Cyrtandra sandwicensis* and *C. grandiflora* occurring on the island Oahu had about 70% fertile pollen grains (Wagner & Herbst, in prep.). In the present investigation hybrids showed high rates of pollen viability, namely between 70 and 99% (see Table 2, Fig. 4D). Even germinated pollen grains with long pollen tubes were found frequently (Fig. 4B). Only one individual of *C. propinqua* \times *C. calpidicarpa* had a highly reduced pollen viability. As abnormal pollen grains generally did not occur more frequently in putative hybrids, reduced pollen viability does not seem to be a reliable criterion for establishing hybrid status.

Especially in late anthesis crystals covering the pollen grain surface and inner wall of the anthers occur occasionally (Fig. 4C, E). Crystals (probably calciumoxalate) have been found in the cells of the connective of the anthers too. No satisfactory answer can be offered for the occurrence of crystals in the pollen sac. Some of them might be sluiced away from the connective cells into the pollen sac by the fixative. An artificial origin of the crystals caused by a reaction between the fixative and some components of the pollen sac content, such as pollenkitt, etc. cannot be excluded (cf. Freytag, 1967).

nerally decreases towards the also occurs in the number of na per 25 μm^2 ranges from 40 (*C. lesii*) to 200 (*C. laxiflora*). It is r within a given species; intra- rtheless, a slight trend can be *ocalyces*, *Apertae* and *Macro-* um^2 at average, rarely 40–60) *e* and *Cylindrocalyces* (40–80 ed are not helpful in supporting

n *Cyrtandra*, SE Asian *Cyrtan-* ting from the Malay Peninsula ting in sculpture and structure: ments lacking, and 2) tectum ractectal elements. The number, it diagnostic features (see Table assessable since the infragenera- a poor state. A very first, inter- ted to the sections *Whitia* and ving supractectal elements and a Pollen morphology might be a awaiian taxa but more data on e statements.

is a widespread phenomenon ; hybrids is the occasionally re- pollen viability by staining the Reduced pollen fertility, how- tween *Cyrtandra sandwicensis* about 70% fertile pollen grains tion hybrids showed high rates e Table 2, Fig. 4D). Even ger- und frequently (Fig. 4B). Only id a highly reduced pollen via- cur more frequently in putative e a reliable criterion for estab-

len grain surface and inner wall tals (probably calciumoxalate) anthers too. No satisfactory an- 1 the pollen sac. Some of them 3 the pollen sac by the fixative. 1 between the fixative and some dtt, etc. cannot be excluded (cf.

ACKNOWLEDGEMENTS

The herbaria at BISH, WU and E kindly provided dried and fixed material for investigation. Dr. M. Kiehn, the National Tropical Botanical Garden (Kauai/Hawaii) and the Bishop Museum (Oahu/Hawaii) are thanked for their support during my field trip on Hawaii. I am grateful to Prof. Dr. M. Hesse and Prof. Dr. A. Weber for critical comments and reading of the manuscript. This study was supported by the Österreichische Nationalbank (Jubiläumsfondsprojekt 3834).

REFERENCES

- Alexander, M.P. 1969. Differential staining of aborted and nonaborted pollen. *Stain Technology* 44: 117–122.
- Clarke, C.B. 1883. *Cyrtandreae* (Gesneracearum tribus). In: A. & C. DeCandolle, *Monogr. Phan.* 5: 1–303.
- Erdtman, G. 1952. Pollen morphology and plant taxonomy. Angiosperms, an introduction to palynology 1. Almqvist & Wiksell, Stockholm.
- Erdtman, G. 1960. The acetolysis method. A revised description. *Svensk Bot. Tidskr.* 54: 561–564.
- Erdtman, G. 1969. Handbook of palynology. Munksgaard, Copenhagen.
- Fægri, K., & J. Iversen. 1975. Textbook of pollen analysis. Munksgaard, Copenhagen.
- Freytag, K. 1967. Über doppelbrechende Lipoide in der Exine und im Pollenöl. *Grana* 7: 3–9.
- Fritze, K., & N.H. Williams. 1988. The taxonomic significance of pollen morphology in the Columnea alliance (Gesneriaceae: Gesnerioideae). *Ann. Missouri Bot. Gard.* 75: 168–191.
- Gerstenberger, P., & P. Leins. 1978. Rasterelektronen-mikroskopische Untersuchung an Blütenknospen von *Physalis philadelphica* (Solanaceae). *Anwendung einer neuen Präparationsmethode. Ber. Dtsch. Bot. Ges.* 91: 381–387.
- Luegmayer, E. 1993. Pollen characters of Old World Gesneriaceae (Cyrtandroideae) with special reference to S.E.-Asian taxa. *Grana* 33 (in press).
- Peterson, R.L., R.E. Hersey & J.D. Brisson. 1978. Embedding softened herbarium material in Spurr's resin for histological studies. *Stain Technology* 53: 1–9.
- Pragowski, J., & W. Punt. 1973. An elucidation of the microreticulate structure of the exine. *Grana* 13: 45–50.
- Roelofs, F.M. 1980. The reproductive biology of *Cyrtandra grandiflora* (Gesneriaceae) on Oahu. *Pacific Sci.* 33: 223–231.
- Spurr, A.R. 1969. A low-viscosity epoxy resin embedding medium for electron microscopy. *J. Ultrastruct. Res.* 26: 31–43.
- St. John, H. 1966. Monograph of *Cyrtandra* (Gesneriaceae) on Oahu, Hawaiian Islands. Bernice P. Bishop Mus. Bull. 229: 1–465.
- Thiéry, J.P. 1967. Mise en évidence des polysaccharides sur coupes fines en microscopie électronique. *J. Microscopie* 6: 987–1018.
- Wagner, W.L., & D.R. Herbst. 1993. Revision of Hawaiian *Cyrtandra* (Gesneriaceae) and a synopsis of hybridization (in prep.).
- Wagner, W.L., D.R. Herbst & S.H. Sohmer. 1990. Manual of the flowering plants of Hawaii. Honolulu, University of Hawaii Press.
- Wodehouse, R.P. 1935. Pollen grains. McGraw-Hill, London, New York.