

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344347055>

# Petrocodon tenuitubus ( Gesneriaceae), a new species from Southeast Yunnan, China

Article · May 2019

---

CITATIONS

0

READS

4

1 author:



Li Chen

Kunming Institute of Botany Chinese Academy of Sciences

6 PUBLICATIONS 1 CITATION

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Taxonomy of Gesneriaceae [View project](#)

DOI: 10.11931/guihaia.gxzw201808033

引文格式: 陈力, 陈文红, 郭世伟, 等. 云南东南部石山苣苔属一新种——细管石山苣苔 [J]. 广西植物, 2019, 39(5): 574–580.  
CHEN L, CHEN WH, GUO SW, et al. *Petrocodon tenuitubus* (Gesneriaceae), a new species from Southeast Yunnan, China [J]. *Guihaia*, 2019, 39(5): 574–580.

## *Petrocodon tenuitubus* (Gesneriaceae), a new species from Southeast Yunnan, China

CHEN Li<sup>1,2</sup>, CHEN Wenhong<sup>1,4</sup>, GUO Shiwei<sup>1,5</sup>, WEN Fang<sup>3,6</sup>, SHUI Yumin<sup>1,4\*</sup>

(1. CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China; 2. School of Life Sciences, Yunnan University, Kunming 650091, China; 3. Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin 541006, Guangxi, China; 4. Karst Conservation Initiative of Yunnan, Kunming 650201, China; 5. University of Chinese Academy of Sciences, Beijing 100049, China; 6. Gesneriad Conservation Center of China, Guilin Botanical Garden, Chinese Academy of Sciences, Guilin 541006, Guangxi, China)

**Abstract:** The newly refined genus *Petrocodon* Hance (Gesneriaceae), mainly distributed from South China to Southwest China, is comprised of 34 species and one variety. A new species, *Petrocodon tenuitubus* W. H. Chen, F. Wen & Y. M. Shui, distributed in the limestone region from Maguan County, Yunnan Province, China, is illustrated and described here. The new species is similar to *P. lui*, *P. hispidus* and *P. jasminiflorus*, but it can be easily distinguished from *P. lui* and *P. hispidus* by its slender and curved corolla tube, three linear to lanceolate bracts and bracteoles, and one disc-shaped stigma. And it also can be easily distinguished from *P. jasminiflorus* by its ovate to orbicular leaves, ovate corolla lobes with obtuse apex, bracts and bracteoles 3, and staminodes 1. The discovery of this new species has certain significance for promoting the excavation of the resources of the Gesneriaceae in limestone area of China. The holotype and paratype specimens are deposited in Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences (KUN); isotype specimens is deposited in Herbarium of Guangxi Institute of Botany (IBK).

**Key words:** flora of Yunnan, limestone cave, new taxon, *Petrocodon*, taxonomy

中图分类号: Q949 文献标识码: A 文章编号: 1000-3142(2019)05-0574-07

## 云南东南部石山苣苔属一新种——细管石山苣苔

陈力<sup>1,2</sup>, 陈文红<sup>1,4</sup>, 郭世伟<sup>1,5</sup>, 温放<sup>3,6</sup>, 税玉民<sup>1,4\*</sup>

(1. 中国科学院昆明植物研究所 东亚植物多样性与生物地理学重点实验室, 昆明 650201; 2. 云南大学 生命科学学院, 昆明 650091; 3. 广西壮族自治区 广西植物研究所, 广西 桂林 541006; 4. 云南喀斯特保护研究会, 昆明 650201; 5. 中国科学院大学, 北京 100049; 6. 中国苦苣苔科植物保育中心, 中国科学院桂林植物园, 广西 桂林 541006)

收稿日期: 2018-10-22

基金项目: 国家自然科学基金(31470306, 31000258, 31860047); 中华人民共和国生态环境部生物多样性调查评估项目; 广西自然科学基金(2017GXNSFAA198006); 广西喀斯特地区植物保护与恢复生态重点实验室基金(17-259-23); 广西科学院科学研究基金(2017YJJ23022) [Supported by the National Natural Science Foundation of China (31470306, 31000258, 31860047); Biodiversity Background Survey and Assessment Program of Ministry of Ecology and Environment of the People's Republic of China; Guangxi Natural Science Foundation Program (2017GXNSFAA198006); Fund of Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain (17-259-23); Science Research Foundation of Guangxi Academy of Sciences (2017YJJ23022)].

作者简介: 陈力(1993-) 男 湖南衡阳人 硕士研究生 主要从事植物系统分类和生物地理学研究 (E-mail) chenlia@mail.kib.ac.cn.

\* 通信作者: 税玉民 博士 研究员 从事热带植物分类学、地理学和生态学研究 (E-mail) ymshui@mail.kib.ac.cn.

摘要: 最近被重新界定的广义石山苣苔属(*Petrocodon* Hance) 是苦苣苔科(Gesneriaceae) 一个中等大小的属,我国目前已知的有 34 种 1 变种,主要分布于我国华南至西南石灰岩地区。该文报道了于云南东南部马关县发现的该属一新种——细管石山苣苔(*Petrocodon tenuitubus* W. H. Chen, F. Wen & Y. M. Shui)。该新种在形态上与陆氏细筒苣苔(*P. lui*)、细筒苣苔(*P. hispidus*) 和长檐苣苔(*P. jasminiflorus*) 相似,但其线形或披针形的苞片和小苞片均为 3 枚,花冠筒细小且弯曲,盘形柱头 1,很容易区别于陆氏细筒苣苔和细筒苣苔;而其叶片卵形至圆形,花冠裂片卵形而尖端钝以及退化雄蕊 3,则显著区别于长檐苣苔。该新种的发现对推进我国石灰岩地区苦苣苔科植物资源的发掘具有一定意义。主模式标本存放于中国科学院昆明植物研究所标本馆(KUN), 等模式标本存放于广西植物研究所标本馆(IBK)。

关键词: 云南植物区系,石灰岩洞穴,新分类群,石山苣苔属,分类学

## 1 Introduction

Genus *Petrocodon* Hance (Gesneriaceae), mainly distributed from South China to Southwest China, is comprised of 34 species and one variety (IPNI 2018). Sixteen species from other eight genera, viz. *Calcareoboea* C. Y. Wu ex H. W. Li, *Didymocarpus* Wall., *Dolicholoma* D. Fang & W. T. Wang, *Lagarosolen* W. T. Wang, *Paralagarosolen* Y. G. Wei, *Primulina* Hance, *Tengia* Chun and *Wentsaiboea* D. Fang & D. H. Qin (Liu et al., 2011; Weber et al., 2011; Clark et al., 2013; Xu et al., 2014; Möller et al., 2016), were merged into this genus and 17 new species of genus *Petrocodon* published in the last more than ten years (Wei, 2007; Jiang et al., 2011; Wen et al., 2012; Chen et al., 2014; Hong et al., 2014; Xu et al., 2014; Li & Wang, 2015; Yu et al., 2015; Guo et al., 2016; Cen et al., 2017; Lu et al., 2017a, 2017b; IPNI, 2018; Zhang et al., 2018).

## 2 Materials and Methods

In October 2016, when the surveys about plant diversity were conducted in Maguan County, Yunnan Province, China, some of authors (YMS, WHC & SWG) collected an unknown species of *Petrocodon* from the limestone area. According to the previous research (Wang, 1984a, 1984b; Pan, 1988; Wang, 1990, 1992; Wang, et al., 1998; Li & Wang, 2004; Wei et al., 2010; Weber et al., 2011; Clark et al., 2013) and the

specimens of KUN, PE, IBK, we confirmed that the species was new to science. And its shapes of corolla and curved tube were slightly similar to *P. lui* (Yan Liu & W. B. Xu) A. Weber & Mich. Möller (Xu et al., 2010; Weber et al., 2011), and its characters of leaf were close to *P. hispidus* (W. T. Wang) A. Weber & Mich. Möller (Wang, 1984b, 1990; Weber et al., 2011), and its corolla tube and stigma were similar to *P. jasminiflorus* (D. Fang et W. T. Wang) A. Weber & Mich. Möller (Wang, 1990; Wang et al., 1998; Weber et al., 2011). But it can be distinguished from *P. lui* by several characters, such as the leaf shape, the number of inflorescences cymes, bracts, bracteoles, and stigma, as well as the indumentum of leaves, inflorescences and flower (Xu et al., 2010), and differs from *P. hispidus* by the number of inflorescences cymes, bracts, bracteoles and stigma, the shape of calyx, corolla and ovary, as well as the indumentum of inflorescences and flower (Wang, 1984b, 1990; Wang et al., 1998), and differs from *P. jasminiflorus* by the leaves characteristics, the shape of corolla, the number of bracts, bracteole and staminodes, as well as the indumentum of leaves, inflorescences and flower. Hence, we describe and illustrate it as a new species, *P. tenuitubus*, endemic to the limestone region in Yunnan Province, Southwest China.

## 3 Taxonomy treatment

***Petrocodon tenuitubus*** W. H. Chen, F. Wen & Y. M. Shui, sp. nov. (Fig. 1–2)

**Type:** China, Yunnan Province: Maguan County, Jinchang Community, 104° 29' 17" E, 22° 46' 29" N, alt. 1 835 m, 29 Oct. 2016, W. H. Chen, S. W. Guo et al., JCL03-011 (holotype, KUN; isotype, IBK).

**Diagnosis:** *P. tenuitubus* is morphologically related to *P. lui*, *P. hispidus* and *P. jasminiflorus*, but it can be easily distinguished from *P. lui* and *P. hispidus* by its slender and curved corolla tube, the number of bracts, bracteole and stigma, as well as the indumentum of inflorescences and flower. And it also can be easily distinguished from *P. jasminiflorus* by its leaves characteristics, the shape of corolla, the number of bracts, bracteole and stami-nodes, as well as the indumentum of leaves, inflorescences and flower.

**Description:** Herbs perennial. Rhizomes 1.0–3.0 cm long, 0.8–1.8 cm in diam. Leaves 5–15, basal; petiolate 0.7–3.5 cm, adaxially green, abaxially reddish-brown, strigose. Leaf blade ovate to orbicular, herbaceous, 2.6–6.8 × 2.0–5.1 cm, obtuse at apex, cordate symmetrically or asymmetrically at base, with crenate margin; adaxially deeply green to yellowish green, densely strigose; abaxially yellowish green to reddish-brown, tomentose; lateral veins pinnate 3–6 pairs, impressed adaxially and prominent abaxially, abaxially bristly. Inflorescences, cymes 1–3, axillary, with 7–15 flowers; peduncles 3.0–6.0 cm long, glandular hairs. Bracts 3, linear to lanceolate, 0.4–0.8 × 0.1–0.3 cm, adaxially tomentose; bracteoles 3, linear to lanceolate, 0.2–0.6 × 0.1–0.2 cm across, adaxially tomentose. Pedicel 0.2–0.6 cm long, densely glandular hairs. Calyx 5-lobed near to the base, lobes linear to lanceolate; lobes equal, 0.3–0.5 × ca. 0.1 cm, densely glandular hairs abaxially. Corolla infundibuliform, pale purple to purple, 1.2–3.6 cm long, throat with a yellow lamina; tube slender, curved, slightly longer than limb, 0.7–1.6 cm long, ca. 0.2 cm in diam. at base and ca. 0.1 cm in the middle, 0.6–0.8 cm in diam. at throat, outside and throat glandular hairs and inside glabrous; limb 5-lobed, adaxial lip 2-lobed, narrowly ovate, apex subacute or obtuse, 0.9–1.1 × 0.5–0.6 cm, abaxial lip 3-lobed, ovate, apex obtuse, 1.2–

1.5 × 0.6–0.8 cm, outside and inside sparsely glandular hairs. Stamens 2, included, adnate at 3–9 mm above the tube base; filament ca. 1 mm long, white, glabrous; anthers oblong, white, glabrous, ca. 1.0 × 0.8 mm, with longitudinal stripe, dorsifixed, confluent apically. Staminodes 3, ca. 0.1 mm long, adnate at 2–8 mm above the corolla tube base. Pistil 4–10 mm long, glandular hairs; ovary ovoid, 1–2 mm long, 1–1.5 mm in diam., glandular hairs; disc ringlike, 0.3 mm; stigma 1, disc-shaped, 0.5–1 mm long. Mature fruit not seen.

**Etymology:** The new species is named by the characteristics of the long and thin corolla tubes.

**Vernacular name:** Xì Guǎn Shí Shān Jù Tái (细管石山苣苔).


**Distribution, habitat and phenology:** The new species, *Petrocodon tenuitubus*, grows under the cliff with only one population in Maguan County of Southeast Yunnan, China. There are some companion species of the new species, such as *Saxifraga mengtzeana* Engl. & Irm-sch., *Mazus pulchellus* Hemsl. ex Forbes & Hemsl., *Aleuritopteris pseudofarinosa* Ching & S. K. Wu. Flowering period occurs from October to November.

**Paratype:** China, Yunnan Province: Maguan County Jinchang Community, 104° 29' 17" E, 22° 46' 29" N, alt. 1 835 m, 10 Nov. 2017, S. W. Guo et al., JCL05-002 (KUN).

## 4 Discussion


The genus *Petrocodon* is endemic to the limestone regions in South China and North Vietnam with habitat fragmentation. Several years ago, two species of *Petrocodon* were discovered in Maguan County and published (Chen et al., 2014). Like species of *Petrocodon*, the three species in Maguan county has the fragment and narrow habitat with a small population (Wang, 1984b, 1990; Wang et al., 1998; Xu et al., 2010; Weber et al., 2011).

The new species grows under the limestone cliff among the non-limestone mountain in the bordering



Note: **A.** Habitat; **B.** Whole plant , with flower; **C.** Leaf adaxial side; **D.** Leaf abaxial side; **E.** Inflorescences; **F.** Bracts and bracteoles; **G.** Flower , front view; **H.** Flower , lateral view; **I.** Opened corolla , showing stamens and pistil; **J.** Stamens ( st.) 2 , confluent apically , and staminodes ( sta.) 3; **K.** Stigma; **L.** Ovary. Photographed by Guo Shiwei ( **A** , **B** , **E** , **G** and **H** ) & Chen Li ( **C** , **D** , **F** , and **I-L** ) .

Fig. 1 **Petrocodon tenuitubus** W. H. Chen , F. Wen & Y. M. Shui



Note: **A-C.** *Petrocodon tenuitubus* **A.** Habit, **B.** Frontal view of flower, **C.** Lateral view of flower. **D-F.** *P. hispidus* **D.** Habit, **E.** Frontal view of flowers, **F.** Lateral view of flowers. **G-I.** *P. lui* **G.** Habit, **H.** Frontal view of flowers, **I.** Lateral view of flowers. **J-L.** *P. jasminiflorus* **J.** Habit, **K.** Frontal view of flowers, **L.** Lateral view of flowers.  
 Photographed by Guo Shiwei (A and C), Wen Fang (B and G-L).

Fig. 2 *Petrocodon tenuitubus* and its morphological related species, *P. hispidus*, *P. lui* and *P. jasminiflorus*

**Table 1** Main different characters among *Petrocodon tenuitubus*, *P. lui*, *P. hispidus* and *P. jasminiflorus*

Character	<i>Petrocodon tenuitubus</i> sp. nov.	<i>Petrocodon lui</i>	<i>Petrocodon hispidus</i>	<i>Petrocodon jasminiflorus</i>
Leaf	Ovate to orbicular	Ovate or broadly ovate, soft	Broadly ovate to orbicular	Narrowly ovate or elliptic
Leaf adaxially	Densely strigose	Glabrous or sparsely puberulent	White appressed hispid	Sparsely white pubescent, margin glandular and ciliate
Leaf abaxially	Tomentose	Puberulent	White appressed hispid	Sparsely white pubescent
Bract	3, linear to lanceolate, tomentose	2, opposite, linear, puberulent	2, opposite, elliptic to oblong, sparsely hispid	1-2, linear, sparsely puberulent
Bracteole	3, linear to lanceolate, tomentose	2, opposite, linear, puberulent	2, opposite, linear, hispidulous and short glandular hairs	1-2, linear, sparsely puberulent
Pedicele	Densely glandular hairs	Puberulent	Hispid	White pubescent and short glandular hairs
Corolla tube	Slender, curved	Slender, curved	Slender, strict	Slender, curved
Corolla	Lobes narrowly ovate to ovate, apex obtuse	Lobes oblong, obovate to suborbicular, apex obtuse	Lobes triangular, apex acute	Lobes narrowly triangular, apex acute
Staminode	3, adnate at 2-8 mm above the corolla tube base	3, adnate at 6-7 cm above the corolla tube base	3, adnate at 10-11.5 mm above the corolla tube base	2, adnate at 4 mm above the corolla tube base
Ovary	Ovoid, glandular hairs	Linear, glandular	Linear, pubescent and few glandular hairs	Narrowly ovoid, pubescent
Stigma	1, disc-shaped	2, slightly split into two parts, aequilateral	2, slightly split into two parts, aequilateral	1, disclike

regions between China and Vietnam. The plants with less than 80 individuals grow on rock faces of the limestone cave near to a road. Because the plants grow beside a new road and their habitat can be easily destroyed. We suggest that it should be classified as “Critically Endangered”: CR B1ab (iii) +2ab (iii), after consulting the IUCN red list categories and criteria (IUCN 2012).

**Acknowledgments** We have so many thanks to Ms. HONG Mengchen and Mr. XI Huihui in Kunming Institute of Botany, Chinese Academy of Sciences for their help in field surveys.

## References:

- CEN HF, FU LF, WEN F, 2017. *Petrocodon urceolatus* sp. nov. (Gesneriaceae) from Zhangjiajie City, northwestern Hunan Province, China [J]. Nord J Bot, 35(3): 334-338.
- CHEN WH, MÖLLER M, SHUI YM, et al., 2014. Three new species of *Petrocodon* (Gesneriaceae), endemic to the limestone areas of Southwest China, and preliminary insights into the diversification patterns of the genus [J]. Syst Bot, 39(1): 316-330.
- CLARK J, WEBER A, MÖLLER M, 2013. A new formal classification of Gesneriaceae [J]. Selbyana, 31(2): 68-94.
- GUO J, MENG T, PANG HB, et al., 2016. *Petrocodon retroflexus* sp. nov. (Gesneriaceae) from a karst cave in Guizhou, China [J]. Nord J Bot, 34(2): 159-164.
- HONG X, ZHOU SB, WEN F, 2014. *Petrocodon villosus* (Gesneriaceae), a new species from Guangxi, China [J]. Blumea, 59(1): 33-36.
- IPNI, 2018. The international plant names index [EB/OL]. [accessed: 30 July 2018]. <http://www.ipni.org/>.
- IUCN, 2012. The IUCN red list of threatened species, Ver. 3.1. 2nd ed [EB/OL]. IUCN Red List Unit, Cambridge U. K. [accessed: 30 July 2018]. <http://www.iucnredlist.org/>.
- JIANG YS, ZHANG Y, WANG Y, et al., 2011. *Petrocodon multiflorus* sp. nov. (Gesneriaceae) from Guangxi, China [J]. Nord J Bot, 29(1): 57-60.
- LI HQ, WANG YQ, 2015. *Petrocodon confertiflorus* (Gesneriaceae), a new species from Guangdong, China [J]. Ann Bot Fenn, 52(1-2): 27-32.
- LI ZY, WANG YZ, 2004. Plants of Gesneriaceae in China [M]. Zhengzhou: Henan Science and Technology Publishing House: 1-721. [李振宇, 王印政, 2004. 中国苦苣苔科植物 [M]. 郑州: 河南科学技术出版社: 1-721.]
- LIU Y, XU WB, HUANG YS, 2011. *Primulina guangxiensis* sp. nov. (Gesneriaceae) from a karst cave in Guangxi, China [J]. Nord J Bot, 29(6): 682-686.

- LU YB, PAN B, HUANG J, et al., 2017a. *Petrocodon pulchri-florus* sp. nov. (Gesneriaceae) from Guangxi, China [J]. *Phytotaxa*, 291(1): 81-88.
- LU YB, HUANG YS, XU WB, et al., 2017b. Floral evolutionary trend of *Petrocodon* (Gesneriaceae) and its significance in classification [J]. *Guihaia*, 37(10): 1227-1239. [卢永彬, 黄俞淞, 许为斌, 等, 2017. 石山苣苔属(苦苣苔科)花形态演化及分类学意义 [J]. *广西植物*, 37(10): 1227-1239.]
- MÖLLER M, WEI YG, WEN F, et al., 2016. You win some you lose some: Updated generic delineations and classification of Gesneriaceae—implications for the family in China [J]. *Guihaia*, 36(1): 44-60. [MÖLLER M, 韦毅刚, 温放, 等, 2016. 得与失: 苦苣苔科新的属级界定与分类系统——中国该科植物之变迁 [J]. *广西植物*, 36(1): 44-60.]
- PAN KY, 1988. Notulae De Gesneriaceis Sinensibus (IX) [J]. *Acta Phytotax Sin*, 26(6): 429-442. [潘开玉, 1988. 中国苦苣苔科的研究(九) [J]. *植物分类学报*, 26(6): 429-442.]
- WANG WT, 1984a. Notulae De Gesneriaceis Sinensibus (V) [J]. *Bull Bot Res*, 4(1): 9-35. [王文采, 1984a. 中国苦苣苔科的研究(五) [J]. *植物研究*, 4(1): 9-35.]
- WANG WT, 1984b. Notulae De Gesneriaceis Sinensibus (VI) [J]. *Acta Bot Yunnan*, 6(1): 11-26. [王文采, 1984b. 中国苦苣苔科的研究(六) [J]. *云南植物研究*, 6(1): 11-26.]
- WANG WT, 1990. Gesneriaceae [M]//WANG WT. *Flora Reipublicae Popularis Sinicae*. Beijing: Science Press, 69: 227, 125-581. [王文采, 1990. 苦苣苔科 [M]//王文采. *中国植物志*. 北京: 科学出版社, 69: 227, 125-581.]
- WANG WT, 1992. Notulae De Gesneriaceis Sinensibus (X) [J]. *Guihaia*, 12(4): 289-300. [王文采, 1992. 中国苦苣苔科的研究(十) [J]. *广西植物*, 12(4): 289-300.]
- WANG WT, PAN KY, LI ZY, et al., 1998. Gesneriaceae [M]//WU CY, RAVEN PH(eds). *Flora of China* (18). Beijing: Science Press; St Louis: Missouri Botanical Garden Press: 244-499.
- WEBER A, WEI YG, PUGLISI C, et al., 2011. A new definition of the genus *Petrocodon* (Gesneriaceae) [J]. *Phytotaxa* 23: 49-67.
- WEI YG, 2007. *Petrocodon ferrugineus* (Gesneriaceae), a new species from Guangxi, China [J]. *Novon*, 17(1): 135-137.
- WEI YG, WEN F, MÖLLER M, et al., 2010. Gesneriaceae of South China [M]. Nanning: Guangxi Science and Technology House: 1-777. [韦毅刚, 温放, MÖLLER M, 等, 2010. 华南苦苣苔科植物 [M]. 南宁: 广西科学技术出版社: 1-777.]
- WEN F, LIANG GY, WEI YG, 2012. *Petrocodon lancifolius* (Gesneriaceae), a new species endemic to a central subtropical zone of Guizhou Province, China [J]. *Phytotaxa*, 49: 45-49.
- XU WB, LIU Y, LIANG YY, et al., 2010. *Lagarosolen lui* Yan Liu et W. B. Xu, a new species of Gesneriaceae from Guangxi, China [J]. *J Wuhan Bot Res*, 28(1): 7-9. [许为斌, 刘演, 梁永延, 等, 2010. 陆氏细筒苣苔——中国广西苦苣苔科一新种 [J]. *武汉植物学研究*, 28(1): 7-9.]
- XU WB, MENG T, ZHANG Q, et al., 2014. *Petrocodon* (Gesneriaceae) in the limestone karsts of Guangxi, China: Three new species and a new Combination based on morphological and molecular evidence [J]. *Syst Bot*, 39(3): 965-974.
- YU XL, LI M, ZHOU JJ, et al., 2015. *Petrocodon hunanensis* (Gesneriaceae), a new species identified by both morphological and molecular evidence from limestone area in Hunan, China [J]. *Phytotaxa*, 195(1): 65-72.
- ZHANG RL, FU LF, LI S, et al., 2018. *Petrocodon asterocalyx*, a new species of Gesneriaceae from Guangxi, China [J]. *Phytotaxa*, 343(3): 259-268.