genes

Article

Characterization and Comparative Analysis of the Complete Mitochondrial Genome of a Limestone-Endemic Endangered Plant Species *Hemiboea yongfuensis* (Gesneriaceae)

Xin-Yue Tao [1,2,3,†], Xin-Mei Qin [2,†], Qiang Zhang [2], Xiao-Li Yang [4], Yong-Bin Lu [2], Yan-Jun Tan [4,5], Peng-Wei Li [2], Xi-Yang Huang [3,*] and Xiang Gan [1,4,5,*]

1 College of Pharmacy, Guilin Medical University, Guilin 541199, China; taoxyyyy@gmail.com
2 Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin 541006, China; xinmeiqin2018@163.com (X.-M.Q.); qiangzhang04@126.com (Q.Z.); yongbinlu@126.com (Y.-B.L.); li_pengweii@126.com (P.-W.L.)
3 Guangxi Key Laboratory of Plant Functional Phytochemicals and Sustainable Utilization, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin 541006, China
4 Guangxi Health Commission Key Laboratory of Disease Proteomics Research, Guilin Medical University, Guilin 541199, China; cncsyxl@126.com (X.-L.Y.); tyj0925@126.com (Y.-J.T.)
5 Scientific Research Center, Guilin Medical University, Guilin 541199, China
* Correspondence: xiyanghuanggxib@126.com (X.-Y.H.); ganxiangdk@163.com (X.G.)
† These authors contributed equally to this work and share the first authorship.

Abstract

Background: *Hemiboea yongfuensis* is a recently discovered critically endangered species. It is exclusive to the limestone regions of Yongfu County, Guilin, Guangxi. Currently, there is a lack of mitogenome data for *Hemiboea* species, hindering the potential of disclosing the evolutionary processes of the mitochondrial genome, which has been far less assembled and shown to be complex in the plant kingdom. Moreover, it prevents potential applications of mitochondrial genome data in phylogenetics and plant adaption, breeding, and conservation. **Results**: In order to reveal the mitochondrial features and variations and explore the usefulness of mitochondrial genes in phylogenetics, in this study, we assembled the complete mitogenome of *H. yongfuensis* using PacBio HiFi long reads, and analyzed its codon usage bias, RNA editing sites, repetitive sequences, sequence lateral transfer, phylogenetic relationships, and synteny. The linear mitochondrial genome assembly we obtained has a length of 619,997 bp and a GC content of 43.63%. The assembly encompasses 61 genes, which include 37 protein-coding genes (PCGs), 21 transfer RNA (tRNA) genes, and 3 ribosomal RNA (rRNA) genes. Importantly, our analysis uncovered a significant presence of repetitive sequences with a high proportion of forward repeats in the mitogenome and significant transposition of sequences from the chloroplast to mitochondrion. Additionally, we revealed the codon usage characteristics of protein-coding genes and identified numerous RNA editing events. Furthermore, we assessed the collinearity of the species in the Gesneriaceae family and found rampant reorganizations. The phylogenetic analyses based on the mitochondrial PCGs for the entire Lamiales order show the monophyly of Gesneriaceae as well as other families and a general high phylogenetic resolution. **Conclusions**: Our study provides the first mitogenome data for *H. yongfuensis* and the genus *Hemiboea*, expanding the rapidly increasing but yet limited plant mitogenome resources. It enhances our understanding of the mitogenome and Lamiales evolution, whereas more potentials of the mitogenome data, such as its possible functions in adaptation to limestone habitats, conservation, and germplasm breeding, remain under-exploited. This first reported *Hemiboea* mitogenome in addition to more mitogenomes from the same and related species would shed further light on these unresolved issues in future studies.

Check for updates

Academic Editor: Bin Yu

Received: 15 December 2025
Revised: 27 January 2026
Accepted: 28 January 2026
Published: 30 January 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

Keywords: organelle genome; repeat sequences; phylogenetics; genomic rearrangement; mitochondrial plastid DNAs

1. Introduction

Mitochondria, the energy-producing centers of cells, are essential for the growth, development, and reproduction of organisms [1]. Research on plant mitochondria started around 1950, coinciding with the first isolation of mitochondria from animal and plant tissues [2]. Plant mitochondria not only participate in many metabolic processes related to energy production and the synthesis and breakdown of different compounds, but also act as carriers of genetic information through their genomes [3]. The mitochondrial genetic system is relatively independent of the cell nucleus, and its genetic information is often inherited from the maternal parent, which somewhat reduces the difficulty of genetic research [4]. However, assembling mitogenomes remains challenging due to their structural diversity, such as branched linear, single linear, circular, and mixed circular–linear forms [5–7]. Traditional short-read sequencing, despite its accuracy, often struggles to resolve these complexities and repetitive sequences, resulting in fragmented assemblies. Long-read sequencing has emerged as a more effective alternative, capable of spanning repetitive regions and accurately determining structural configurations. Additionally, the presence of extensive non-coding sequences, high-frequency repetitions, RNA editing, and plastid genome insertions [8–11] further complicates the assembly process. Some of these features also lead to substantial differences in the size of plant mitogenomes [12]. To date, there are significantly fewer complete plant mitogenomes reported compared to plant chloroplast genomes [10]. Although plant mitogenomes vary greatly in structure and size, the mitochondrial protein-coding genes tend to have conserved features. Some of these genes have been utilized in phylogenetic studies of plants, showing potential and importance in plant phylogenetics [13,14].

The Gesneriaceae family includes around 160 genera and more than 3800 species, making it one of the larger groups of tropical plants [15]. The family includes herbs, vines, and shrubs, which often have ornamental value because of their attractive flowers. Certain species have been utilized in traditional medicine mainly for addressing issues such as fever, cough, the common cold, snake bites, and pain, along with various infectious and inflammatory conditions [16]. The Gesneriaceae family, a member of the Lamiales order, provides important insights into evolutionary history and mechanisms of Lamiales featured with zygomorphic and bilabiate flowers. However, assembling plant mitogenomes has been a longstanding conundrum due to sequencing technique limitations that the short reads produced by Sanger sequencing or next-generation sequencing are unable to span the long repeats frequently occurring in plant mitogenomes, along with a lack of efficient assembling tools for dealing with such complex plant mitogenomes. In addition, the usefulness of the mitochondrial sequences, for example, in phylogenetics, has not been adequately tested or realized, further preventing the endeavor of studying plant mitogenomes. Hence, despite being a large family, only four species have had their mitogenomes assembled to date: *Boea hygrometrica* and *Haberlea rhodopensis*, which are known for extreme desiccation tolerance; *Primulina hunanensis*, an endangered species adapted to cave environments; and *Oreocharis esquirolii*, a species with rare actinomorphic corolla in the family [5,17–19]. Fortunately, the fast developments of PacBio and Oxford Nanopore long-read sequencing techniques, along with the newly designed assembling tools based on long-read data in the recent past, have dramatically improved the accessibility of the complete plant mitogenomes [20–22].

Hemiboea C. B. Clarke (1888) is a genus of perennial herbaceous plants in Gesneriaceae, consisting of about 42 species and five varieties [23]. They are mainly found in southern China, with some species also present in northern Vietnam and Japan [24,25]. Among these, *H. yongfuensis* has been found only in Yongfu County, Guilin, Guangxi, growing on limestone rocks (Figure 1). Owing to its limited population size, restricted distribution area, and the risk of habitat degradation, the species was assessed as critically endangered according to the IUCN standards [26]. Previous phylogenetic studies of *Hemiboea* were based solely on plastid and nuclear ribosomal ITS sequences [27]. As of now, no mitogenome of the genus *Hemiboea* has been assembled or utilized for phylogenetic studies, despite the availability of complete mitogenomes for 53 other species within the order Lamiales in NCBI.




Figure 1. *Hemiboea yongfuensis* in the limestone habitat, North Guangxi, China. Photograph by Yongbin Lu.

In this study, we assembled the complete mitogenome of *H. yongfuensis* by PacBio long-read sequencing, which represents the first assembled mitogenome in the genus. We comprehensively analyzed its gene composition, codon usage preferences, repetitive sequences, RNA editing, and homology between the mitochondrial and chloroplast genomes, as well as synteny with the previously published mitogenomes from the Gesneriaceae family and phylogenetic relationships in Lamiales. We mainly aimed to uncover the evolutionary processes of the mitogenome and its usefulness in the phylogenetics in Lamiales. Although the focus was on the mitogenome, the chloroplast genome of the same individual was also assembled as it serves as a critical reference for identifying plastid-derived DNA insertions in the mitogenome. The assembled mitogenome expands the genomic resources of the Gesneriaceae family, a diverse plant group with substantial ornamental, horticultural, as well as medicinal values, which would serve as a valuable asset for developing molec-

ular markers to further explore the evolution and adaptation and inform conservation and breeding.

2. Materials and Methods

2.1. Plant Materials, DNA Extraction, and Sequencing

Although *H. yongfuensis* was assessed as critically endangered (CR) according to the IUCN standards [26], it has not been included in any (officially) issued list of protected plant species. The individuals we collected were from a wild, non-private area (i.e., not from any nature reserve or owned by any third party), and therefore no collection permit or license was required. Fresh leaves were collected from a wild individual in Luojin Town, Yongfu County, Guilin city, Guangxi, southern China (110°8′17″ E, 24°58′36″ N, 284 m). The fresh leaves were rinsed with ultrapure water and immediately frozen in liquid nitrogen. Total DNA was then extracted using a modified CTAB method [28], and its concentration and integrity were assessed with NanoDrop spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) and Qubit fluorometer (Thermo Fisher Scientific, Waltham, MA, USA). For short-read sequencing, genomic DNA was randomly fragmented using a Covaris ultrasonic homogenizer to generate a short-read library with 150 bp inserts. Paired-end sequencing was conducted using the Illumina platform NovaSeq 6000 platform, yielding about 218 Gb of sequencing data. For long-read sequencing, a PacBio HiFi long-read library (SMRTbell library) with an average insert size of about 17 kb was prepared. Sequencing was conducted on the PacBio Revio platform, producing 49 Gb of sequencing data with an average read length of 17,269 bp and an N50 value of 16,876 bp. All library preparation and sequencing were carried out by Wuhan Benagen Technology Company (Wuhan, China).

2.2. Assembly and Annotation of Mitogenome

Mitogenome assembly was performed using PacBio HiFi long-read sequencing data. The raw sequencing data were assembled de novo with Flye (v2.9.1-b1780) [29] under default parameters to generate graphical assembly outputs in GFA (Graphical Fragment Assembly) format. All resultant contigs in FASTA format were subjected to BLASTn (v2.13.0) [30] analysis with the reference mitogenomes of *Arabidopsis thaliana* and *P. hunanensis* to identify contigs that include mitogenome sequences. A customized BLAST database was constructed using makeblastdb, followed by homology searches implemented with the parameters -evalue 1×10^{-5} -outfmt 6 -max_hsps 10 -word_size 7 -task blastn-short. The GFA file was visualized and analyzed using Bandage software (v0.8.1) [31], and mitochondrial contigs were filtered based on BLASTn results to obtain the draft mitogenome of *H. yongfuensis* (Figure 2A). Each node in the visualization corresponds to an assembled contig, and a black line connecting two nodes indicates overlapping regions between the respective contig sequences. Collectively, these sequences form a complex, multi-branched circular genome structure. For key branching nodes, connections supported by longer sequencing reads are prioritized as follows: long reads were mapped to the sequences at these branching nodes, and a long read that continuously aligns with two connected sequences along the black line serves as evidence for their connection. When multiple alternative connections exist at a branching node, those supported by a greater number of long reads are given priority. Through this process, the most plausible genome structure (Figure 2B) of the *H. yongfuensis* mitogenome was ultimately determined.

Annotation of the mitogenomes was performed using PMGA [32] and Geseq software (v2.03) [33], with *A. thaliana* (NC_037304), *P. hunanensis* (NC_087815), and *O. esquirolii* (PQ850635) used as references. After further manual verification and correction, the annotated mitogenomes have been deposited in GenBank. The mitogenome annotation results were visualized by OGDRAW [34].




Figure 2. Assembly results of *H. yongfuensis* mitogenome. (**A**) Schematic of *H. yongfuensis* mitogenome assembly; (**B**) main conformation of *H. yongfuensis* mitogenome.

2.3. Assembly and Annotation of Chloroplast Genome

The chloroplast genome assembly was conducted using GetOrganelle software (v1.7.5) [35] with default parameters from short-read sequencing data. The chloroplast genome was annotated using the CPGAVAS2 web service (http://www.herbalgenomics.org/cpgavas2/ accessed on 17 September 2025) [36] with *H. yongfuensis* (GenBank: NC_079573) as the reference. Finally, the assembled and annotated *H. yongfuensis* chloroplast genome was submitted to GenBank. This assembly and annotation primarily aimed to facilitate chloroplast–mitochondrial homology analysis using the same sample.

2.4. Analysis of Codon Usage and Repeated Sequences

The protein-coding genes (PCGs) of the mitogenome were initially extracted using PhyloSuite software (v1.2.2) [37]. Thereafter, CodonW software (v1.4.2) [38] was utilized to calculate the relative synonymous codon usage (RSCU) values and conduct codon usage bias analysis.

Simple sequence repeats (SSRs) were detected using MISA (v2.1) (https://webblast.ipk-gatersleben.de/misa/ accessed on 29 June 2025) [39] with minimum repeat unit thresholds set to 10, 5, 4, 3, 3, and 3 for mono- to hexa-nucleotide motifs, respectively. Tandem repeat analysis was performed using Tandem Repeats Finder (TRF v4.09; https://tandem.bu.edu/trf/trf.unix.help.html accessed on 29 June 2025) [40] with default settings. For dispersed repeats, Reputer (https://bibiserv.cebitec.uni-bielefeld.de/reputer/ accessed on 29 June 2025) [41] was employed to identify four categories: forward (F), palindromic (P), reverse (R), and complementary (C) repeats.

2.5. Chloroplast–Mitochondrial Homology Analysis

The homologous sequence analysis between the mitochondrial and plastid genomes (MTPTs) was conducted with BLASTN (v2.13.0) [30] under the following settings: -evalue 1×10^{-5}, -word_size 10, and -outfmt 6. Finally, TBtools (v2.309) [42] was employed to visualize chloroplast-to-mitochondrial gene transfer events.

2.6. RNA Editing Sites Prediction

RNA editing sites within all PCGs encoded by the *H. yongfuensis* mitogenome were predicted using Deepred-Mt [43], a convolutional neural network (CNN)-based model. Predictions with probability scores exceeding 0.9 were retained for subsequent analysis.

2.7. The Mitogenome Comparative Analyses of H. yongfuensis to Other Gesneriaceae

The mitogenome sequences of the five Gesneriaceae species including *H. yongfuensis* underwent pairwise BLAST (v2.16.0) comparisons. Searches were conducted with an E-value threshold of 1×10^{-5} and a word size of 7. Resulting alignments were retained only if they exhibited over 80% nucleotide identity and spanned more than 500 base pairs, with the output format specified as -outfmt 6. Subsequently, the multiple synteny plot was generated using TBtools software (v2.309).

Gene nucleotide sequences were aligned using MAFFT (v7.505) [44]. Nucleotide diversity (Pi) for each aligned gene was then calculated in DnaSP (v5.10) [45]. The resulting Pi values were visualized as a line graph using GraphPad Prism (v10.1.2).

2.8. Phylogenetic Analysis

Mitogenome sequences of 56 species, encompassing all the available Lamiales species and 2 Solanaceae species designated as the outgroup, were retrieved from NCBI. A total of 25 conserved PCGs from these species were extracted using PhyloSuite (v1.1.16) [37]. These PCGs were subsequently aligned with MAFFT (v7.505) [44] and segments that were excessively divergent or poorly aligned were masked using the alignmentFilter package through a sliding window grouping–regrouping strategy (with the stringent parameter *prob* set to 0.0001) [46]. The resulting alignment was then employed to construct a maximum likelihood (ML) phylogenetic tree using IQ-TREE (v1.6.12) [47,48], with branch support assessed through 1000 ultrafast bootstrap replicates (-B 1000) and 1000 SH-aLRT tests (-alrt 1000). The phylogenetic tree was visualized using ITOL (v6) [49].

3. Results

3.1. Structural Characteristics of the Mitogenome of H. yongfuensis

We successfully assembled the mitogenome of *H. yongfuensis* and presented it in a specific linear structure, which, however, may be one of its possible conformations. Figure 2A displays the schematic of *H. yongfuensis* mitogenome assembly, including nodes with their respective length and sequencing depth (Table S1). The sequences of the linear contig produced after resolving the multifurcated nodes (at the black connecting lines) using long-read data are illustrated in Figure 2B, with a specific solution path of edge_3-edge_2-edge_1.

The mitogenome of *H. yongfuensis* measures a total length of 619,997 bp, with a GC content of 43.63% (Figure 3). The annotation identified 37 distinct PCGs, consisting of 24 core mitochondrial genes and 13 non-core genes, along with 21 tRNA genes (including 5 multi-copy tRNAs) and 3 rRNA genes (Table 1). Among the core genes, there is one protein transport subunit gene (*mtt*B), nine NADH dehydrogenase genes (*nad*1, *nad*2, *nad*3, *nad*4, *nad*4L, *nad*5, *nad*6, *nad*7, and *nad*9), and one gene for ubiquinol–cytochrome c reductase (*cob*). Additionally, the core gene set includes three cytochrome c oxidase genes (*cox*1, *cox*2, and *cox*3), one maturase gene (*mat*R), and four cytochrome c biogenesis genes (*ccm*FC, *ccm*FN, *ccm*C, and *ccm*B), along with five ATP synthase genes (*atp*1, *atp*4, *atp*6, *atp*8, and *atp*9). Non-core genes comprise four ribosomal large subunit genes (*rpl*2, *rpl*5, *rpl*10, *rpl*16), seven ribosomal small subunit genes (*rps*3, *rps*4, *rps*7, *rps*10, *rps*12, *rps*13, *rps*14), and two succinate dehydrogenase genes (*sdh*3, *sdh*4).

Table 1. The mitogenome encoding genes of *H. yongfuensis*.

Group of Genes	Name of Genes
ATP synthase	*atp*1, *atp*4, *atp*6, *atp*8, *atp*9
NADH dehydrogenase	*nad*1, *nad*2, *nad*3, *nad*4, *nad*4L, *nad*5, *nad*6, *nad*7, *nad*9
Cytochrome *b*	*cob*
Cytochrome *c* biogenesis	*ccm*B, *ccm*C, *ccm*FC, *ccm*FN
Cytochrome *c* oxidase	*cox*1, *cox*2, *cox*3
Maturases	*mat*R
Protein transport subunit	*mtt*B
Ribosomal protein large subunit	*rpl*2, *rpl*5, *rpl*10, *rpl*16
Ribosomal protein small subunit	*rps*3, *rps*4, *rps*7, *rps*10, *rps*12, *rps*13, *rps*14
Succinate dehydrogenase	*sdh*3, *sdh*4
Ribosome RNA	*rrn*5, *rrn*18, *rrn*26
Transfer RNA	*trn*A-UGC, *trn*C-GCA, *trn*D-GUC, *trn*E-UUC (×2), *trn*F-GAA (×2), *trn*fM-CAU (×3), *trn*G-GCC, *trn*H-GUG, *trn*I-CAU, *trn*K-UUU, *trn*L-CAA (×2), *trn*L-UAA, *trn*M-CAU, *trn*N-GUU, *trn*P-UGG (×2), *trn*Q-UUG, *trn*S-GGA, *trn*S-GCU, *trn*S-UGA, *trn*W-CCA, *trn*Y-GUA

Note: The numerical value enclosed in parentheses signifies the number of copies of the gene, such as (×2) representing the existence of two copies.




Figure 3. The mitogenome map of *H. yongfuensis* with annotation.

3.2. Codon Usage of PCGs

In the complete mitogenome of *H. yongfuensis*, we identified 10,546 codons among the 37 PCGs detected. The mitogenome of *H. yongfuensis* encodes 21 different amino acids, utilizing a total of 64 distinct codons. The analysis of codon usage bias, as shown in Figure 4, revealed that aside from the start codon ATG, tryptophan (Trp, TGG), and alanine (Ala, GCA), three of which have RSCU values of one, there exists a general preference in codon usage among the mitochondrial PCGs. Specifically, 31 codons exhibit RSCU values greater than one, indicating a preferential usage for those amino acids. For instance, alanine (Ala) demonstrates a strong preference for the codon GCT, boasting the highest RSCU value of 1.59, while the stop codon TAG has the lowest frequency at 0.43.




Figure 4. Codon usage of 21 amino acids in the mitogenome PCGs of *H. yongfuensis*. The *x*-axis indicates the codon family, while the *y*-axis displays the frequency of use. Different colors are used to identify codons that correspond to the same amino acid.

3.3. Analysis of Repeat Sequences

An analysis of repetitive sequences within the mitochondrial genome. of *H. yongfuensis* revealed a total of 121 simple sequence repeats (SSRs), as detailed in Table S2. This includes 27 monomeric SSRs (22.31%), 28 dimeric SSRs (23.14%), 17 trimeric SSRs (14.05%), and 43 tetrameric SSRs (35.53%), which represented the largest proportion. There are also six pentameric SSRs (4.96%), but no hexameric SSRs were found (Figure 5A). Furthermore, four tandem repeat sequences, each with a matching degree greater than 79% and lengths varying from 16 to 24 bp, were identified in the mitogenome (Figure 5B; Table S3). The mitogenome of *H. yongfuensis* also contains a significant number of dispersed repeats, totaling 294 pairs, with lengths of 30 bp or more (Table S4). This consists of 122 pairs of palindromic repeats (P), 171 pairs of forward repeats (F), and 1 pair of reverse repeats (R), with no complementary repeats detected (Figure 5B; Table S4). The longest palindromic repeat measures 141 bp, while the longest forward repeat extends to 4556 bp (Table S4).




Figure 5. Histogram of repetitive sequence analyses. (**A**) The horizontal axis represents SSR types, while the vertical axis shows the number of repetitive fragments. (**B**) The horizontal axis shows the repeat types, while the vertical axis illustrates the number of repetitive fragments.

3.4. Homology Analysis of Genomic Sequences

A sequence similarity analysis revealed 74 homologous mitochondrial plastid sequences (MTPTs) in *H. yongfuensis*. These sequences with over 80% similarity between the cpDNA and mitogenome measure a total of 72,759 bp, which accounts for 11.74% of the overall mitogenome length (Figure 6). Of these, MTPT1 is the longest, with a length of 4956 bp (Table S5).

The annotation of these homologous sequences identified a total of 31 complete genes among the 74 MTPTs analyzed. This includes 21 PCGs such as *rps*11, *rps*14, *rps*8, *rps*2, *ndh*J, *ycf*15, *psb*B, *rpl*23, *rpl*2, *rpl*14, *rpl*36, *psb*A, *psb*C, *psb*D, *psb*L, *psb*F, *psb*E, *pet*L, *pet*G, *inf* A,

and *rpo*B, as well as 10 tRNA genes: *trn*A-UGC, *trn*D-GUC, *trn*H-GUG, *trn*I-CAU, *trn*L-CAA, *trn*L-UAA, *trn*N-GUU, *trn*W-CCA, *trn*S-GGA, and *trn*fM-CAU. In addition, several incomplete chloroplast genes were also detected within these homologous sequences, including *ycf*2, *ndh*K, *rpo*A, *rps*4, and others (Table S5).




Figure 6. Homology analysis of the mitochondrial and chloroplast genomes of *H. yongfuensis*. The blue arc represents the mitogenome, the green arc represents the chloroplast genome, and orange lines between the arcs correspond to homologous sequence fragments.

3.5. RNA Editing Site Prediction

Deepred-mt software (https://github.com/aedera/deepredmt/ accessed on 20 September 2025) predicted RNA editing events in the 37 PCGs from the mitogenome of *H. yongfuensis*. With a predictive performance score of 0.9 (where 0 indicates no editing and 1 indicates editing, with values closer to 1 suggesting a higher likelihood of editing), a total of 429 potential RNA editing sites were identified across these 37 PCGs. These sites were consistently characterized by C to U conversions (Table S6). Notably, the *ccm*B gene had the highest number of RNA editing sites, with 37 identified. Following this, the *nad4* gene experienced 36 RNA editing events. In comparison, the *rps*7 and *atp*1 genes

each had only one editing site (Figure 7). Additionally, further analysis revealed that a total of 44 codon variants were associated with RNA editing sites. Among these variants, it was predicted that most amino acids would change from hydrophilic to hydrophobic, accounting for approximately 47.09%. Meanwhile, 43.12% of the amino acids were expected to maintain their hydrophobicity or hydrophilicity, and 9.32% were predicted to change from hydrophobic to hydrophilic. Notably, the study identified that the codons may have been edited into stop codons in *atp*6 and *rps*10 genes (Table S7).




Figure 7. Number of RNA editing sites predicted for the 37 PCGs in the *H. yongfuensis* mitogenome.

3.6. Collinearity Analysis

In the collinearity analysis, any collinear blocks shorter than 0.5 kb and with similarity below 80% were omitted. The results indicated that a significant number of homologous collinear blocks were identified between *H. yongfuensis* and its allies within Gesneriaceae (Figure 8). A comparative analysis was conducted on the mitogenomes of between *H. yongfuensis* and *P. hunanensis*, and between *H. yongfuensis* and *O. esquirolii*. It was revealed that the highly homologous regions in the mitogenomes exhibit a broader distribution range between *H. yongfuensis* and *P. hunanensis* (Figure 8). Notably, the length of the homologous region in *H. yongfuensis* is 447,434 bp, accounting for 72.17% of the total length of its mitogenome. In *P. hunanensis*, the homologous region spans 447,862 bp, accounting for 77.86% of the total length of its mitogenome (Table S8). The largest collinear block was in comparison with *P. hunanensis*, spanning approximately 13,188 bp (Table S8). The arrangement of collinear blocks within the mitogenomes of the four species in Gesneriaceae varied significantly, and that *H. yongfuensis* exhibited extensive genomic rearrangements (Figure 8) against its allies from the same family.




Figure 8. Collinearity analysis of *H. yongfuensis* and its allies in Gesneriaceae. The red arcs denote reverse homologous regions, and the gray arcs indicate positive homologous regions.

3.7. Nucleotide Diversity

Nucleotide diversity (Pi) is a key metric used to evaluate genetic variation in nucleotide sequences among different species and populations. Areas with high variability can serve as potential molecular markers for differentiating between populations. An analysis of Pi was carried out on mitochondrial genes from the four Gesneriaceae species. The results revealed that the mitochondrial gene with the highest variability was *cox*2 (Pi = 0.03333), followed by *nad*2 (Pi = 0.025) (Figure 9). This finding suggests that the nucleotide sequences of mitochondrial PCGs in Gesneriaceae are highly conserved.




Figure 9. Nucleotide diversity of the mitochondrial genes from the five Gesneriaceae species. The horizontal axis represents gene names, and the vertical axis represents the Pi values of the genes.

3.8. Phylogenetic Analyses

To investigate the phylogenetic significance of mitochondrial genes and the position of *H. yongfuensis* within Lamiales, a phylogenetic tree was constructed using DNA sequences from the 25 PCGs from 54 Lamiales species whose mitochondrial genomes had been publicly available. Two species from the Solanales order were used as outgroups. The selected PCGs included *atp*1, *atp*4, *atp*6, *atp*8, *atp*9, *ccm*B, *ccm*C, *cob*, *cox*1, *cox*2, *cox*3, *mat*R, *mtt*B, *nad*1, *nad*2, *nad*3, *nad*4, *nad*4L, *nad*5, *nad*6, *nad*7, *nad*9, *rps*3, and *rps*12. The results (Figure 10) indicate that all the eleven families are monophyletic in the order. Remarkably, the monophyly of Gesneriaceae is strongly supported (BS = 100), with *H. yongfuensis*

embedded within this family. Within Gesneriaceae, the phylogenetic relationships are as follows: *H. yongfuensis* is closely related to *O. esquirolii* (BS = 50); this clade is further grouped with *P. hunanensis* to form a larger lineage (BS = 95), which is then sister to *B. hygrometrica* (BS = 99), and the entire clade encompassing these four taxa ultimately forms a sister relationship with *Hab. rhodopensis* (BS = 100).




Figure 10. Phylogenetic position of *H. yongfuensis* within Lamiales. The maximum likelihood tree is based on nucleotide sequences of 25 conserved mitochondrial protein-coding genes selected for phylogenetic analysis. It includes 54 species from Lamiales, with *Solanum melongena* and *Nicotiana longiflora* set as outgroups. Numbers at branch nodes indicate bootstrap support values. Branches are colored according to their respective plant families. GenBank accession numbers are provided after each of the species names. Phylogenetic analysis was conducted using MAFFT for sequence alignment and IQ-TREE for tree construction.

4. Discussion

Currently, complete mitogenomes data for Gesneriaceae plants remain extremely limited, with only four species, *B. hygrometrica*, *Hab. rhodopensis*, *P. hunanensis*, and *O. esquirolii* [5,17–19], having had their mitogenomes completely determined. To comprehensively understand the diversity and evolutionary characteristics of Gesneriaceae mi-

togenomes, we sequenced, assembled, and conducted systematic analysis of the complete mitochondrial genome of *H. yongfuensis*. Notably, the assembled *H. yongfuensis* mitogenome exhibits a linear structure, representing the first complete mitogenome obtained within the genus *Hemiboea*. Comparisons between *H. yongfuensis* and previously reported Gesneriaceae mitogenomes reveal that despite significant variations in genome size across genera, from 619,997 bp (*H. yongfuensis*) to 454,871 bp (*O. esquirolii*), the GC content remains highly consistent (ranging from 43.54% in *P. hunanensis* to 43.8% in *O. esquirolii*). This supports the hypothesis of a relatively stable GC content in higher plant genomes during evolution [12]. Furthermore, the reported Gesneriaceae species encompass both linear structures (*H. yongfuensis*, *P. hunanensis*, and *O. esquirolii*) and circular structures (*B. hygrometrica* and *Hab. rhodopensis*), further corroborating the view that plant mitogenome structures exhibit high dynamism and complexity.

Despite the significant variations observed in the size and structure of plant mitogenomes, the number of mitochondrial genes among terrestrial plants has remained relatively stable [50,51]. We further evaluated the conservation of the *H. yongfuensis* mitogenome at the gene composition level. Recent annotation results indicate that the mitogenome of *H. yongfuensis* comprises 61 genes, which include 37 PCGs, 21 tRNAs, and 3 rRNAs. Among these genes, 37 PCGs are commonly shared by most angiosperms [6]. This relative stability in gene content provides a basis for examining other features that contribute to mitogenome diversity. Differences in the frequencies of synonymous codon usage can offer valuable supplementary data for phylogenetic analyses [52]. Our investigation into the codon usage preferences in 37 PCGs revealed that the same amino acid can be encoded by as many as six possible codons (such as leucine, serine, and arginine). Moreover, we observed distinct differences in the usage frequencies of codons associated with each amino acid. For those codons exhibiting a synonymous codon usage index greater than 1, a clear preference for adenine and thymine (AT) bases at the third codon position is evident, a trend that is commonly found in higher plants [53].

The migration of chloroplast sequences to the mitogenome is a common phenomenon observed in higher plants [54–58]. These transferred sequences, known as mitochondrial plastid DNA transfers (MTPTs), had been integrated into the mitogenome and contribute to its complexity and diversity. *H. yongfuensis* also exhibits this chloroplast sequence transfer, which is consistent with findings from other species in Gesneriaceae [5]. Specifically, the total length of the transferred segments within the mitogenome of *H. yongfuensis* measures 72,759 bp, which constitutes 11.74% of the entire mitogenome length, which slightly exceeds the commonly reported values ranging from 3% to 11.5% [59,60]. Previous research indicates that genes migrating from chloroplasts to mitochondria often become pseudogenes over time, losing their functional significance due to processes such as sequence recombination [61]. Interestingly, the MTPT fragments identified in *H. yongfuensis* contain multiple intact PCGs and tRNA genes. The integrity of the sequence may indicate that the transfer event is relatively recent or has been somewhat conserved throughout evolution, suggesting that it may still have functional significance. However, more evidence is needed to determine whether it is expressed and performs a function. Additionally, the underlying mechanisms that facilitate intergenomic sequence migration remain to be further investigated [62].

In addition to intergenomic transfer, transcript-level modification may also contribute to mitochondrial functional variation [63]. RNA editing, the process that primarily consists of the conversion of cytosine (C) to uracil (U) in mitochondrial gene transcripts, impacts protein function and structure and thus plays a crucial role in evolution [62,64]. In our study of the *H. yongfuensis* mitogenome, 429 RNA editing sites were detected across its 37 PCGs. Among the Gesneriaceae species, this count is greater than that reported for *Hab.*

rhodopensis (419 sites), yet it is lower than *P. hunanensis* (455 sites) [5,17,18]. In *H. yongfuensis*, it is predicted that 47.09% of the amino acids in PCGs will change from hydrophilic to hydrophobic due to RNA editing. These changes assist in protein folding and the formation of secondary structures, which help optimize protein conformation and function [65,66]. The biological significance of RNA editing extends beyond gene expression regulation; it also plays a crucial role in the adaptive evolution of plants. Research indicates that the rate of non-synonymous substitutions due to C-to-U editing events is higher than that of synonymous substitutions. Moreover, these editing sites show greater conservation at the DNA level, suggesting that they have an adaptive role in natural selection [67]. Additionally, RNA editing is closely linked to plant development, the regulation of flowering time, and responses to environmental stresses [62]. In stressful environments, such as limestone areas, plants require a stable energy supply and effective stress responses. This adaptability is determined by a combination of multiple mechanisms [68–70]. *H. yongfuensis* is adapted to the limestone habitat and has flowering period divergence compared to its known sister species *Hem. subcapitata* from the same areas [71]. However, whether RNA editing has driven the divergence and adaptation of the species merits further investigation.

Plant mitogenomes contain many repetitive sequences, such as simple, tandem, and scattered repeats [72–74], which affect their size and structure. These repeats can influence genome evolution and diversity by enabling recombination and gene transfer. Previous studies have demonstrated that mitogenome rearrangements exhibit high dynamicity across different lineages and may also be influenced by evolutionary processes. For instance, in *Brassica* plants, mitogenome rearrangements are associated with evolutionary selection during domestication and breeding processes, suggesting they may be linked to adaptation under environmental stress [75]. In the mitogenome of *H. yongfuensis*, the simple repeat sequences identified are mainly monomers and dimers, predominantly consisting of A/T repeats and AT/TA repeat structures. This feature is consistent with two other similar species in the Gesneriaceae family [5,19] and aligns with a general trend observed in most terrestrial plants [76–78].

In angiosperms, mitogenome data have proven to be very effective at clarifying deep relationships, serving as a valuable resource for phylogenetic and evolutionary studies of these plants [79,80]. In this study, we utilized mitogenomes to determine the phylogenetic position of *H. yongfuensis* in Gesneriaceae as well as in Lamiales. The results show that *B. hygrometrica*, *Hab. rhodopensis*, *P. hunanensis*, *O. esquirolii*, and *H. yongfuensis* form a highly supported clade, with a bootstrap support (BS) of 100. Notably, *H. yongfuensis* is more closely related to *O. esquirolii*, which is consistent with the findings of the previous studies based on chloroplast genomes [81]. However, the bootstrap support for this relationship is relatively low (BS = 50), necessitating further verification using nuclear genes. Additionally, due to the imbalance between plant mitochondrial and chloroplast data, more comprehensive sampling of mitogenomes is also required for better resolution of the phylogeny. Furthermore, the results showed somewhat significant variation in nucleotide diversity (Pi) among the mitochondrial PCGs of *H. yongfuensis*, ranging from complete conservativeness in *atp*9 (Pi = 0) to higher variability in *cox*2 (Pi = 0.033) and nad2 (Pi = 0.025). This variation may reflect different selective pressures resulting from functional constraints and evolutionary processes. However, adaptive interpretations remain uncertain without more extensive sampling and integrated analyses that explicitly link genomic variation to ecological differences.

5. Conclusions

In this study, we successfully assembled and annotated the complete mitogenome of *H. yongfuensis*, marking the first report of a complete mitogenome for the genus *Hemiboea*.

The mitogenome features a linear structure, measuring 619,997 bp in total, and including 37 PCGs, 21 tRNA genes, and 3 rRNA genes. The study revealed genomic rearrangements and frequent plastid-derived DNA insertions, whereas the gene content, GC composition, and codon usage patterns are largely preserved. While the taxonomic sampling in this study was somewhat limited, the findings indicate that mitochondrial data can provide valuable phylogenetic insights. Overall, these results contribute to the limited mitochondrial DNA database of Gesneriaceae plants, enhance our understanding of their evolution, and provide important genetic data for the taxonomic classification, systematic evolution, and species conservation.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/genes17020167/s1. Supplementary Material S1. Table S1: results based on Flye assembly; Table S2: SSRs in the mitochondrial genome of *H. yongfuensis*; Table S3: tandem repeat sequences in the mitochondrial genome of *H. yongfuensis*; Table S4: dispersed repeat sequences in the mitochondrial genome of *H. yongfuensis*; Table S5: the homologous DNA fragment in the *H. yongfuensis* mitochondrial genome; Table S6: potential RNA editing sites in PCGs of the *H. yongfuensis* mitochondrial genome; Table S7: amino acid changes in potential RNA editing sites of PCGs in the *H. yongfuensis* mitochondrial genome; Table S8: homologous co-linear blocks of *H. yongfuensis* with closely related species.

Author Contributions: X.G. and X.-Y.H. designed the study. X.-Y.T., X.-M.Q., and Y.-B.L. performed the data analyses. X.-Y.T. and X.-M.Q. drafted the manuscript. X.-L.Y., Y.-J.T., and P.-W.L. collected the plant materials and data. X.-M.Q., Q.Z., X.G., and X.-Y.H. revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by the National Natural Science Foundation of China (No. 32460057), Guangxi Natural Science Foundation (No. GuikeAD23026329, 2023GXNSFAA026307, and 2020GXNSFAA297202), and National Key Research and Development Program of China (No. 2022YFD1600302).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The complete mitochondrial and chloroplast genome sequences of *H. yongfuensis* are available at GenBank with accession numbers: PX394567 and PX388148.

Conflicts of Interest: The authors declare that they have no conflicts of interest.

References

1. Sun, M.Y.; Zhang, M.Y.; Chen, X.N.; Liu, Y.Y.; Liu, B.B.; Li, J.M.; Wang, R.Z.; Zhao, K.J.; Wu, J. Rearrangement and domestication as drivers of Rosaceae mitogenome plasticity. *BMC Biol.* **2022**, *20*, 181. [CrossRef] [PubMed]
2. Møller, I.M.; Rasmusson, A.G.; Van, A.O. Plant mitochondria—Past, present and future. *Plant J.* **2021**, *108*, 912–959. [CrossRef] [PubMed]
3. Cheng, Y.; He, X.X.; Priyadarshani, S.V.G.N.; Wang, Y.; Ye, L.; Shi, C.; Ye, K.Z.; Zhou, Q.; Luo, Z.Q.; Deng, F.; et al. Assembly and comparative analysis of the complete mitochondrial genome of *Suaeda glauca*. *BMC Genom.* **2021**, *22*, 167. [CrossRef] [PubMed]
4. Wallace, D.C.; Singh, G.; Lott, M.T.; Hodge, J.A.; Schurr, T.G.; Lezza, A.M.S.; Elsas, L.J.; Nikoskelainen, E.K. Mitochondrial DNA mutation associated with Leber's hereditary optic neuropathy. *Science* **1988**, *242*, 1427–1430. [CrossRef]
5. Chen, L.L.; Dong, X.; Huang, H.; Xu, H.X.; Rono, P.C.; Cai, X.Z.; Hu, G.W. Assembly and comparative analysis of the initial complete mitochondrial genome of *Primulina hunanensis* (Gesneriaceae): A cave-dwelling endangered plant. *BMC Genom.* **2024**, *25*, 322. [CrossRef]
6. Wang, Y.; Chen, S.J.; Chen, J.J.; Chen, C.J.; Lin, X.J.; Peng, H.; Zhao, Q.; Wang, X.Y. Characterization and phylogenetic analysis of the complete mitochondrial genome sequence of *Photinia serratifolia*. *Sci. Rep.* **2023**, *13*, 770. [CrossRef]
7. Kozik, A.; Rowan, B.A.; Lavelle, D.; Berke, L.; Schranz, M.E.; Michelmore, R.W.; Christensen, A.C. The alternative reality of plant mitochondrial DNA: One ring does not rule them all. *PLoS Genet.* **2019**, *15*, e1008373. [CrossRef]
8. Cole, L.W.; Guo, W.H.; Mower, J.P.; Palmer, J.D. High and variable rates of repeat-mediated mitochondrial genome rearrangement in a genus of plants. *Mol. Biol. Evol.* **2018**, *35*, 2773–2785. [CrossRef]

9. Li, C.; Liu, H.; Qin, M.; Tan, Y.J.; Ou, X.L.; Chen, X.Y.; Wei, Y.; Zhang, Z.J.; Lei, M. RNA editing events and expression profiles of mitochondrial protein-coding genes in the endemic and endangered medicinal plant, *Corydalis saxicola*. *Front. Plant Sci.* **2024**, *15*, 1332460. [CrossRef]
10. Wang, J.; Kan, S.L.; Liao, X.Z.; Zhou, J.W.; Tembrock, L.R.; Daniell, H.; Jin, S.X.; Wu, Z.Q. Plant organellar genomes: Much done, much more to do. *Trends Plant Sci.* **2024**, *29*, 754–769. [CrossRef]
11. Wu, Z.Q.; Liao, X.Z.; Zhang, X.N.; Tembrock, L.R.; Broz, A. Genomic architectural variation of plant mitochondria—A review of multichromosomal structuring. *J. Syst. Evol.* **2022**, *60*, 160–168. [CrossRef]
12. Sloan, D.B.; Alverson, A.J.; Chuckalovcak, J.P.; Wu, M.; McCauley, D.E.; Palmer, J.D.; Taylor, D.R. Rapid evolution of enormous, multichromosomal genomes in flowering plant mitochondria with exceptionally high mutation rates. *PLoS Biol.* **2012**, *10*, e1001241. [CrossRef] [PubMed]
13. Song, N.; Liang, A.P.; Bu, C.P. A molecular phylogeny of Hemiptera inferred from mitochondrial genome sequences. *PLoS ONE* **2012**, *7*, e48778. [CrossRef] [PubMed]
14. Qiu, Y.L.; Li, L.B.; Wang, B.; Xue, J.Y.; Hendry, T.A.; Li, R.Q.; Brown, J.W.; Liu, Y.; Hudson, G.T.; Chen, Z.D. Angiosperm phylogeny inferred from sequences of four mitochondrial genes. *J. Syst. Evol.* **2010**, *48*, 391–425. [CrossRef]
15. Ogutcen, E.; Christe, C.; Nishii, K.; Salamin, N.; Möller, M.; Perret, M. Phylogenomics of Gesneriaceae using targeted capture of nuclear genes. *Mol. Phylogenet. Evol.* **2021**, *157*, 107068. [CrossRef]
16. Verdan, M.H.; Stefanello, M.É.A. Secondary metabolites and biological properties of Gesneriaceae species. *Chem. Biodivers.* **2012**, *9*, 2701–2731. [CrossRef]
17. Zhang, T.W.; Fang, Y.J.; Wang, X.M.; Deng, X.; Zhang, X.W.; Hu, S.N.; Yu, J. The complete chloroplast and mitochondrial genome sequences of *Boea hygrometrica*: Insights into the evolution of plant organellar genomes. *PLoS ONE* **2012**, *7*, e30531. [CrossRef]
18. Baev, V.; Ivanova, Z.; Yahubyan, G.; Toneva, V.; Apostolova, E.; Minkov, G.; Minkov, I. Analysis of the complete mitochondrial genome sequence of the resurrection plant *Haberlea rhodopensis*. *Acta Biochim. Pol.* **2021**, *68*, 277–286. [CrossRef]
19. Chen, L.; Yan, R.R.; Yang, C.Y.; Ling, L.Z.; Bai, X.X.; Ren, Q.F.; Hu, G.X. Mitochondrial genome analysis of the endangered *Oreocharis esquirolii*: Insights into evolutionary adaptation and conservation. *BMC Plant Biol.* **2025**, *25*, 827. [CrossRef]
20. Rhoads, A.; Au, K.F. PacBio sequencing and its applications. *Genom. Proteom. Bioinform.* **2015**, *13*, 278–289. [CrossRef]
21. Brown, C.G.; Clarke, J. Nanopore development at Oxford Nanopore. *Nat. Biotechnol.* **2016**, *34*, 810–811. [CrossRef] [PubMed]
22. Bi, C.; Shen, F.; Han, F.; Qu, Y.; Hou, J.; Xu, K.; Xu, L.-A.; He, W.; Wu, Z.; Yin, T. PMAT: An efficient plant mitogenome assembly toolkit using low-coverage HiFi sequencing data. *Hortic. Res.* **2024**, *11*, uhae023. [CrossRef] [PubMed]
23. Feng, M.; Zhang, J.S. The complete chloroplast genome of *Hemiboea subacaulis* var. *jiangxiensis* Z. Y. Li 1983 (Gesneriaceae), an endemic species in China. *Mitochondrial DNA Part B* **2024**, *9*, 1497–1500. [CrossRef] [PubMed]
24. Weber, A. Gesneriaceae and Scrophulariaceae: Robert Brown and now. *Telopea* **2004**, *10*, 543–571.
25. Li, Z.Y.; Wang, Y.Z. *Plants of Gesneriaceae in China*; Henan Science and Technology Publishing House: Zhengzhou, China, 2005.
26. Huang, Z.P.; Li, J.H.; Pan, B.; Qin, X.M.; Zhang, Q.; Lu, Y.B. *Hemiboea yongfuensis* (Gesneriaceae): A cryptic and critically endangered new species from North Guangxi, China. *Nord. J. Bot.* **2020**, *38*, e02435. [CrossRef]
27. Cui, Y.F.; Zhou, P.; Xiang, K.L.; Zhang, Q.; Yan, H.; Zhang, L.G.; Pan, B.; Huang, Y.S.; Guo, Z.Y.; Li, Z.Y.; et al. Plastome evolution and phylogenomics of Trichosporeae (Gesneriaceae) with its morphological characters appraisal. *Front. Plant Sci.* **2023**, *14*, 1160535. [CrossRef]
28. Porebski, S.; Bailey, L.G.; Baum, B.R. Modification of a CTAB DNA extraction protocol for plants containing high polysaccharide and polyphenol components. *Plant Mol. Biol. Report.* **1997**, *15*, 8–15. [CrossRef]
29. Kolmogorov, M.; Yuan, J.; Lin, Y.; Pevzner, P.A. Assembly of long, error-prone reads using repeat graphs. *Nat. Biotechnol.* **2019**, *37*, 540–546. [CrossRef]
30. Chen, Y.; Ye, W.C.; Zhang, Y.D.; Xu, Y.S. High speed BLASTN: An accelerated MegaBLAST search tool. *Nucleic Acids Res.* **2015**, *43*, 7762–7768. [CrossRef]
31. Wick, R.R.; Schultz, M.B.; Zobel, J.; Holt, K.E. Bandage: Interactive visualization of de novo genome assemblies. *Bioinformatics* **2015**, *31*, 3350–3352. [CrossRef]
32. Li, J.L.; Ni, Y.; Lu, Q.Q.; Chen, H.M.; Liu, C. PMGA: A plant mitochondrial genome annotator. *Plant Commun.* **2025**, *6*, 101191. [CrossRef] [PubMed]
33. Tillich, M.; Lehwark, P.; Pellizzer, T.; Ulbricht-Jones, E.S.; Fischer, A.; Bock, R.; Greiner, S. GeSeq—Versatile and accurate annotation of organelle genomes. *Nucleic Acids Res.* **2017**, *45*, W6–W11. [CrossRef] [PubMed]
34. Greiner, S.; Lehwark, P.; Bock, R. OrganellarGenomeDRAW (OGDRAW) version 1.3.1: Expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* **2019**, *47*, W59–W64. [CrossRef] [PubMed]
35. Jin, J.J.; Yu, W.B.; Yang, J.B.; Song, Y.; dePamphilis, C.W.; Yi, T.S.; Li, D.Z. GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biol.* **2020**, *21*, 241. [CrossRef]
36. Shi, L.C.; Chen, H.M.; Jiang, M.; Wang, L.Q.; Wu, X.; Huang, L.F.; Liu, C. CPGAVAS2, an integrated plastome sequence annotator and analyzer. *Nucleic Acids Res.* **2019**, *47*, W65–W73. [CrossRef]

37. Zhang, D.; Gao, F.L.; Jakovlić, I.; Zou, H.; Zhang, J.; Li, W.X.; Wang, G.T. PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Mol. Ecol. Resour.* **2020**, *20*, 348–355. [CrossRef]
38. Iriarte, A.; Lamolle, G.; Musto, H. Codon usage bias: An endless tale. *J. Mol. Evol.* **2021**, *89*, 589–593. [CrossRef]
39. Beier, S.; Thiel, T.; Münch, T.; Scholz, U.; Mascher, M. MISA-web: A web server for microsatellite prediction. *Bioinformatics* **2017**, *33*, 2583–2585. [CrossRef]
40. Benson, G. Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Res.* **1999**, *27*, 573–580. [CrossRef]
41. Kurtz, S. REPuter: The manifold applications of repeat analysis on a genomic scale. *Nucleic Acids Res.* **2001**, *29*, 4633–4642. [CrossRef]
42. Chen, C.J.; Chen, H.; Zhang, Y.; Thomas, H.R.; Frank, M.H.; He, Y.H.; Xia, R. TBtools: An integrative toolkit developed for interactive analyses of big biological data. *Mol. Plant* **2020**, *13*, 1194–1202. [CrossRef]
43. Edera, A.A.; Small, I.; Milone, D.H.; Sanchez-Puerta, M.V. Deepred-Mt: Deep representation learning for predicting C-to-U RNA editing in plant mitochondria. *Comput. Biol. Med.* **2021**, *136*, 104682. [CrossRef] [PubMed]
44. Katoh, K.; Standley, D.M. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol. Biol. Evol.* **2013**, *30*, 772–780. [CrossRef] [PubMed]
45. Librado, P.; Rozas, J. DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **2009**, *25*, 1451–1452. [CrossRef] [PubMed]
46. Zhang, Q.; Qin, X.M.; Lu, Y.B.; Li, P.W.; Huang, X.Y. A Comprehensive Alignment-Filtering Methodology Improves Phylogeny Particularly by Filtering Overly Divergent Segments. Available online: https://www.biorxiv.org/content/10.1101/2023.12.26.573321v1.full.pdf (accessed on 1 May 2025).
47. Huelsenbeck, J.P.; Ronquist, F. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **2001**, *17*, 754–755. [CrossRef]
48. Nguyen, L.T.; Schmidt, H.A.; Von, H.A.; Minh, B.Q. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **2015**, *32*, 268–274. [CrossRef]
49. Letunic, I.; Bork, P. Interactive Tree of Life (iTOL) v4: Recent updates and new developments. *Nucleic Acids Res.* **2019**, *47*, W256–W259. [CrossRef]
50. Wang, L.; Liu, X.; Wang, Y.D.; Ming, X.J.; Qi, J.S.; Zhou, Y.Q. Comparative analysis of the mitochondrial genomes of four *Dendrobium* species (Orchidaceae) reveals heterogeneity in structure, synteny, intercellular gene transfer, and RNA editing. *Front. Plant Sci.* **2024**, *15*, 1429545. [CrossRef]
51. Guo, W.H.; Grewe, F.; Fan, W.S.; Young, G.J.; Knoop, V.; Palmer, J.D.; Mower, J.P. *Ginkgo* and *Welwitschia* mitogenomes reveal extreme contrasts in gymnosperm mitochondrial evolution. *Mol. Biol. Evol.* **2016**, *33*, 1448–1460. [CrossRef]
52. Zhang, Y.; Shen, Z.N.; Meng, X.R.; Zhang, L.M.; Liu, Z.G.; Liu, M.J.; Zhang, F.; Zhao, J. Codon usage patterns across seven Rosales species. *BMC Plant Biol.* **2022**, *22*, 65. [CrossRef]
53. Yang, H.X.; Li, W.H.; Yu, X.L.; Zhang, X.Y.; Zhang, Z.Y.; Liu, Y.X.; Wang, W.X.; Tian, X.X. Insights into molecular structure, genome evolution and phylogenetic implication through mitochondrial genome sequence of *Gleditsia sinensis*. *Sci. Rep.* **2021**, *11*, 14850. [CrossRef] [PubMed]
54. Zhang, T.T.; Liu, H.; Gao, Q.Y.; Yang, T.; Liu, J.N.; Ma, X.F.; Li, Z.H. Gene transfer and nucleotide sequence evolution by *Gossypium* cytoplasmic genomes indicates novel evolutionary characteristics. *Plant Cell Rep.* **2020**, *39*, 765–777. [CrossRef] [PubMed]
55. Chen, J.F.; Yu, R.X.; Dai, J.H.; Liu, Y.; Zhou, R.C. The loss of photosynthesis pathway and genomic locations of the lost plastid genes in a holoparasitic plant *Aeginetia indica*. *BMC Plant Biol.* **2020**, *20*, 199. [CrossRef] [PubMed]
56. Cui, H.N.; Ding, Z.; Zhu, Q.L.; Wu, Y.; Qiu, B.Y.; Gao, P. Comparative analysis of nuclear, chloroplast, and mitochondrial genomes of watermelon and melon provides evidence of gene transfer. *Sci. Rep.* **2021**, *11*, 1595. [CrossRef]
57. Niu, Y.F.; Gao, C.W.; Liu, J. Mitochondrial genome variation and intergenomic sequence transfers in *Hevea* species. *Front. Plant Sci.* **2024**, *15*, 1234643. [CrossRef]
58. Wang, S.Y.; Qiu, J.; Sun, N.; Han, F.C.; Wang, Z.F.; Yang, Y.; Bi, C.W. Characterization and comparative analysis of the first mitochondrial genome of *Michelia* (Magnoliaceae). *Genom. Commun.* **2025**, *2*, e001. [CrossRef]
59. Alverson, A.J.; Wei, X.; Rice, D.W.; Stern, D.B.; Barry, K.; Palmer, J.D. Insights into the evolution of mitochondrial genome size from complete sequences of *Citrullus lanatus* and *Cucurbita pepo* (Cucurbitaceae). *Mol. Biol. Evol.* **2010**, *27*, 1436–1448. [CrossRef]
60. Yurina, N.P.; Odintsova, M.S. Mitochondrial genome structure of photosynthetic eukaryotes. *Biochemistry* **2016**, *81*, 101–113. [CrossRef]
61. Anderson, B.M.; Krause, K.; Petersen, G. Mitochondrial genomes of two parasitic *Cuscuta* species lack clear evidence of horizontal gene transfer and retain unusually fragmented *ccmFC* genes. *BMC Genom.* **2021**, *22*, 816. [CrossRef]
62. Tang, W.; Luo, C. Molecular and functional diversity of RNA editing in plant mitochondria. *Mol. Biotechnol.* **2018**, *60*, 935–945. [CrossRef]
63. Wang, S.; Wang, H.; Xu, Z.; Jiang, S.; Shi, Y.; Xie, H.; Wang, S.; Hua, J.; Wu, Y. m^6A mRNA modification promotes chilling tolerance and modulates gene translation efficiency in *Arabidopsis*. *Plant Physiol.* **2023**, *192*, 1466–1482. [CrossRef] [PubMed]

64. Lo, G.C.; Pesole, G.; Picardi, E. REDIdb 3.0: A comprehensive collection of RNA editing events in plant organellar genomes. *Front. Plant Sci.* **2018**, *9*, 482. [CrossRef] [PubMed]
65. Li, Z.C.; Liu, Z.Q.; Zhong, W.Q.; Huang, M.H.; Wu, N.; Xie, Y.; Dai, Z.; Zou, X.Y. Large-scale identification of human protein function using topological features of interaction network. *Sci. Rep.* **2016**, *6*, 37179. [CrossRef] [PubMed]
66. Brenner, W.G.; Mader, M.; Müller, N.A.; Hoenicka, H.; Schroeder, H.; Zorn, I.; Fladung, M.; Kersten, B. High level of conservation of mitochondrial RNA editing sites among four *Populus* species. *G3 Genes Genomes Genet.* **2019**, *9*, 709–717. [CrossRef]
67. Chu, D.; Wei, L. The chloroplast and mitochondrial C-to-U RNA editing in *Arabidopsis thaliana* shows signals of adaptation. *Plant Direct* **2019**, *3*, e00169. [CrossRef]
68. Tang, S.; Liu, J.; Lambers, H.; Zhang, L.; Liu, Z.; Lin, Y.; Kuang, Y. Increase in leaf organic acids to enhance adaptability of dominant plant species in karst habitats. *Ecol. Evol.* **2021**, *11*, 10277–10289. [CrossRef]
69. Sand-Jensen, K.; Hammer, K.J. Moss cushions facilitate water and nutrient supply for plant species on bare limestone pavements. *Oecologia* **2012**, *170*, 305–312. [CrossRef]
70. Feng, C.; Wang, J.; Wu, L.; Kong, H.; Yang, L.; Feng, C.; Wang, K.; Rausher, M.; Kang, M. The genome of a cave plant, *Primulina huaijiensis*, provides insights into adaptation to limestone karst habitats. *New Phytol.* **2020**, *227*, 1249–1263. [CrossRef]
71. Huang, J.; Xiang, X.G.; Lu, Y.B.; Pan, B.; Zhang, Q. *Hemiboea pterocaulis* comb. et stat. nov. (Gesneriaceae), a new critically endangered species segregated from *H. subcapitata*. *Nord. J. Bot.* **2018**, *36*, e01468. [CrossRef]
72. Dong, S.S.; Zhao, C.X.; Chen, F.; Liu, Y.H.; Zhang, S.Z.; Wu, H.; Zhang, L.S.; Liu, Y. The complete mitochondrial genome of the early flowering plant *Nymphaea colorata* is highly repetitive with low recombination. *BMC Genom.* **2018**, *19*, 614. [CrossRef]
73. Guo, W.H.; Zhu, A.D.; Fan, W.S.; Mower, J.P. Complete mitochondrial genomes from the ferns *Ophioglossum californicum* and *Psilotum nudum* are highly repetitive with the largest organellar introns. *New Phytol.* **2017**, *213*, 391–403. [CrossRef] [PubMed]
74. Gandini, C.L.; Sanchez-Puerta, M.V. Foreign plastid sequences in plant mitochondria are frequently acquired via mitochondrion-to-mitochondrion horizontal transfer. *Sci. Rep.* **2017**, *7*, 43402. [CrossRef] [PubMed]
75. Yang, J.; Liu, G.; Zhao, N.; Chen, S.; Liu, D.; Ma, W.; Hu, Z.; Zhang, M. Comparative mitochondrial genome analysis reveals the evolutionary rearrangement mechanism in *Brassica*. *Plant Biol.* **2016**, *18*, 527–536. [CrossRef] [PubMed]
76. Wang, R.; Luo, Y.J.; Lan, Z.; Qiu, D.S. Insights into structure, codon usage, repeats, and RNA editing of the complete mitochondrial genome of *Perilla frutescens* (Lamiaceae). *Sci. Rep.* **2024**, *14*, 13940. [CrossRef]
77. Guo, Y.Y.; Yang, J.X.; Li, H.K.; Zhao, H.S. Chloroplast genomes of two species of *Cypripedium*: Expanded genome size and proliferation of AT-biased repeat sequences. *Front. Plant Sci.* **2021**, *12*, 609729. [CrossRef]
78. Bi, C.W.; Lu, N.; Xu, Y.Q.; He, C.P.; Lu, Z.H. Characterization and analysis of the mitochondrial genome of common bean (*Phaseolus vulgaris*) by comparative genomic approaches. *Int. J. Mol. Sci.* **2020**, *21*, 3778. [CrossRef]
79. Lin, D.L.; Shao, B.Y.; Gao, Z.Y.; Li, J.W.; Li, Z.H.; Li, T.Y.; Huang, W.C.; Zhong, X.; Xu, C.; Chase, M.W.; et al. Phylogenomics of angiosperms based on mitochondrial genes: Insights into deep node relationships. *BMC Biol.* **2025**, *23*, 45.
80. Soe, T.; Kong, J.L.; Xu, B.; Kan, J.H.; Tembrock, L.R.; Nie, L.Y.; Jia, M.L. Mitochondrial genomes of four *Dendrobium* species (Orchidaceae): A comprehensive analysis of structural diversity, synteny, and plastid-derived sequences. *Genom. Commun.* **2025**, *2*, e018. [CrossRef]
81. Zimmer, E.A.; Roalson, E.H.; Skog, L.E.; Boggan, J.K.; Idnurm, A. Phylogenetic relationships in the Gesnerioideae (Gesneriaceae) based on nrDNA ITS and cpDNA trn L-F and trn E-T spacer region sequences. *Am. J. Bot.* **2002**, *89*, 296–311. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.